

**FHIT Antibody (monoclonal) (M03)**

Mouse monoclonal antibody raised against a partial recombinant FHIT.

Catalog # AT2046a

**Specification**

---

**FHIT Antibody (monoclonal) (M03) - Product Information**

Application	WB, E
Primary Accession	<a href="#">P49789</a>
Other Accession	<a href="#">BC032336</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	16858

**FHIT Antibody (monoclonal) (M03) - Additional Information**

Gene ID 2272

**Other Names**

Bis(5'-adenosyl)-triphosphatase, AP3A hydrolase, AP3Aase, Diadenosine 5', 5'''-P1, P3-triphosphate hydrolase, Dinucleosidetriphosphatase, Fragile histidine triad protein, FHIT

**Target/Specificity**

FHIT (AAH32336, 31 a.a. ~ 130 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

FHIT Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


**FHIT Antibody (monoclonal) (M03) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**FHIT Antibody (monoclonal) (M03) - Images**



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



FHIT monoclonal antibody (M01), clone 1E3 Western Blot analysis of FHIT expression in HL-60 (Cat # AT2046a )



Detection limit for recombinant GST tagged FHIT is approximately 0.03ng/ml as a capture antibody.

**FHIT Antibody (monoclonal) (M03) - Background**

This gene, a member of the histidine triad gene family, encodes a diadenosine 5',5'''-P1,P3-triphosphate hydrolase involved in purine metabolism. The gene encompasses the common fragile site FRA3B on chromosome 3, where carcinogen-induced damage can lead to translocations and aberrant transcripts of this gene. In fact, aberrant transcripts from this gene have been found in about half of all esophageal, stomach, and colon carcinomas. Alternatively spliced transcript variants have been found for this gene.

### **FHIT Antibody (monoclonal) (M03) - References**

Homozygous deletion but not mutation of exons 5 and 8 of the fragile histidine triad (FHIT) gene is associated with features of differentiated thyroid carcinoma. Yin DT, et al. *Ann Clin Lab Sci*, 2010 Summer. PMID 20689140. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. *Diabetes Care*, 2010 Jul 13. PMID 20628086. [The relationship between FHIT and WWOX expression and clinicopathological features in hepatocellular carcinoma] Lin J, et al. *Zhonghua Gan Zang Bing Za Zhi*, 2010 May. PMID 20510001. [Genetic and epigenetic changes of FHIT gene in patients with esophageal cancer] Tsao I, et al. *Tsitol Genet*, 2009 Nov-Dec. PMID 20458975. Genomic profiling of adult acute lymphoblastic leukemia by single nucleotide polymorphism oligonucleotide microarray and comparison to pediatric acute lymphoblastic leukemia. Okamoto R, et al. *Haematologica*, 2010 Sep. PMID 20435627.