

HOXD8 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant HOXD8.

Catalog # AT2430a

Specification

HOXD8 Antibody (monoclonal) (M03) - Product Information

Application	WB
Primary Accession	P13378
Other Accession	NM_019558
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	31911

HOXD8 Antibody (monoclonal) (M03) - Additional Information**Gene ID** 3234**Other Names**

Homeobox protein Hox-D8, Homeobox protein Hox-4E, Homeobox protein Hox-54, HOXD8, HOX4E

Target/Specificity

HOXD8 (NP_062458, 126 a.a. ~ 190 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

HOXD8 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


HOXD8 Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HOXD8 Antibody (monoclonal) (M03) - Images



HOXD8 monoclonal antibody (M03), clone 5E11 Western Blot analysis of HOXD8 expression in HepG2 (Cat # L019V1).

HOXD8 Antibody (monoclonal) (M03) - Background

This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conserved family of transcription factors that play an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, located on different chromosomes, consisting of 9 to 11 genes arranged in tandem. This gene is one of several homeobox HOXD genes located in a cluster on chromosome 2. Deletions that remove the entire HOXD gene cluster or the 5' end of this cluster have been associated with severe limb and genital abnormalities. In addition to effects during embryogenesis, this particular gene may also play a role in adult urogenital tract function.

HOXD8 Antibody (monoclonal) (M03) - References

Altered transmission of HOX and apoptotic SNPs identify a potential common pathway for clubfoot. Ester AR, et al. Am J Med Genet A, 2009 Dec. PMID 19938081. Identification of targets of Prox1 during in vitro vascular differentiation from embryonic stem cells: functional roles of HoxD8 in lymphangiogenesis. Harada K, et al. J Cell Sci, 2009 Nov 1. PMID 19825936. High-density association study of 383 candidate genes for volumetric BMD at the femoral neck and lumbar spine among older men. Yerges LM, et al. J Bone Miner Res, 2009 Dec. PMID 19453261. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Limb malformations and the human HOX genes. Goodman FR. Am J Med Genet, 2002 Oct 15. PMID 12357469.