

## **IDH2 Antibody (monoclonal) (M01)**

Mouse monoclonal antibody raised against a partial recombinant IDH2.

Catalog # AT2479a

### **Specification**

---

#### **IDH2 Antibody (monoclonal) (M01) - Product Information**

Application	WB, IHC, E
Primary Accession	<a href="#">P48735</a>
Other Accession	<a href="#">NM_002168</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	50909

#### **IDH2 Antibody (monoclonal) (M01) - Additional Information**

**Gene ID** 3418

##### **Other Names**

Isocitrate dehydrogenase [NADP], mitochondrial, IDH, ICD-M, IDP, NADP(+)-specific ICDH, Oxalosuccinate decarboxylase, IDH2

##### **Target/Specificity**

IDH2 (NP\_002159, 354 a.a. ~ 451 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

##### **Dilution**

WB~~1:500~1000

##### **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

##### **Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

##### **Precautions**

IDH2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


#### **IDH2 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### IDH2 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.52 KDa) .




Western Blot analysis of IDH2 expression in transfected 293T cell line by IDH2 monoclonal antibody (M01), clone 5F11.

Lane 1: IDH2 transfected lysate(50.9 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to IDH2 on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged IDH2 is approximately 0.03ng/ml as a capture antibody.

### **IDH2 Antibody (monoclonal) (M01) - Background**

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex.

### **IDH2 Antibody (monoclonal) (M01) - References**

1. Altered global methylation and hydroxymethylation status in vulvar lichen sclerosus - further support for epigenetic mechanisms. Gambichler T, Terras S, Kreuter A, Skrygan M Br J Dermatol. 2013 Oct 27. doi: 10.1111/bjd.12702.