

IRF5 Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant IRF5.

Catalog # AT2548a

Specification

IRF5 Antibody (monoclonal) (M05) - Product Information

Application	IF, WB, IHC
Primary Accession	Q13568
Other Accession	NM_002200
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	56044

IRF5 Antibody (monoclonal) (M05) - Additional Information**Gene ID** 3663**Other Names**

Interferon regulatory factor 5, IRF-5, IRF5

Target/Specificity

IRF5 (NP_002191, 395 a.a. ~ 504 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

IRF5 Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.


IRF5 Antibody (monoclonal) (M05) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IRF5 Antibody (monoclonal) (M05) - Images




Immunofluorescence of monoclonal antibody to IRF5 on A-431 cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



IRF5 monoclonal antibody (M05), clone 2D4 Western Blot analysis of IRF5 expression in A-431 (Cat # AT2548a)



Immunoperoxidase of monoclonal antibody to IRF5 on formalin-fixed paraffin-embedded human testis. [antibody concentration 0.8 ug/ml]

IRF5 Antibody (monoclonal) (M05) - Background

This gene encodes a member of the interferon regulatory factor (IRF) family, a group of transcription factors with diverse roles, including virus-mediated activation of interferon, and modulation of cell growth, differentiation, apoptosis, and immune system activity. Members of the IRF family are characterized by a conserved N-terminal DNA-binding domain containing tryptophan (W) repeats. Multiple transcript variants encoding different isoforms have been found for this gene, and a 30-nt indel polymorphism (SNP rs60344245) can result in loss of a 10-aa segment.

IRF5 Antibody (monoclonal) (M05) - References

Genome-wide meta-analyses identify three loci associated with primary biliary cirrhosis. Liu X, et al. Nat Genet, 2010 Aug. PMID 20639880. Variants at IRF5-TNPO3, 17q12-21 and MMEL1 are associated with primary biliary cirrhosis. Hirschfield GM, et al. Nat Genet, 2010 Aug. PMID 20639879. Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Genetic Differences between Five European Populations. Moskvina V, et al. Hum Hered, 2010 Jul 8. PMID 20616560. Dengue hemorrhagic fever is associated with polymorphisms in JAK1. Silva LK, et al. Eur J Hum Genet, 2010 Jun 30. PMID 20588308.