

LG MN Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant LG MN.

Catalog # AT2703a

Specification

LG MN Antibody (monoclonal) (M02) - Product Information

Application	E
Primary Accession	O99538
Other Accession	BC003061
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	49411

LG MN Antibody (monoclonal) (M02) - Additional Information

Gene ID 5641

Other Names

Legumain, Asparaginyl endopeptidase, Protease, cysteine 1, LG MN, PRSC1

Target/Specificity

LG MN (AAH03061.1, 1 a.a. ~ 433 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

LG MN Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


LG MN Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

LGMN Antibody (monoclonal) (M02) - Images



Detection limit for recombinant GST tagged LGMN is approximately 0.1ng/ml as a capture antibody.

LGMN Antibody (monoclonal) (M02) - Background

This gene encodes a cysteine protease that has a strict specificity for hydrolysis of asparaginyl bonds. This enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and appears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A fully mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. Overexpression of this gene may be associated with the majority of solid tumor types. This gene has a pseudogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two variants encode the same isoform.

LGMN Antibody (monoclonal) (M02) - References

1.Targeting Cell Surface Alpha(v)beta(3,5) Integrins Increases Therapeutic Efficacies of a Legumain Protease-Activated Auristatin Prodrug.Liu Y, Bajjuri KM, Liu C, Sinha SC.Mol Pharm. 2011 Oct 31.