

MVD Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant MVD.

Catalog # AT2940a

Specification

MVD Antibody (monoclonal) (M01) - Product Information

Application	IF, WB, E
Primary Accession	P53602
Other Accession	NM_002461
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	43405

MVD Antibody (monoclonal) (M01) - Additional Information

Gene ID 4597

Other Names

Diphosphomevalonate decarboxylase, Mevalonate (diphospho)decarboxylase, MDDase, Mevalonate pyrophosphate decarboxylase, MVD, MPD

Target/Specificity

MVD (NP_002452, 301 a.a. ~ 398 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MVD Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


MVD Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MVD Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to MVD on A-431 cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.52 KDa) .




MVD monoclonal antibody (M01), clone 2A7 Western Blot analysis of MVD expression in A-431 (

(Cat # AT2940a)



Western Blot analysis of MVD expression in transfected 293T cell line by MVD monoclonal antibody (M01), clone 2A7.

Lane 1: MVD transfected lysate(43.4 KDa).
Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged MVD is approximately 0.1ng/ml as a capture antibody.

MVD Antibody (monoclonal) (M01) - Background

The enzyme mevalonate pyrophosphate decarboxylase catalyzes the conversion of mevalonate pyrophosphate into isopentenyl pyrophosphate in one of the early steps in cholesterol biosynthesis. It decarboxylates and dehydrates its substrate while hydrolyzing ATP.

MVD Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. Human mevalonate diphosphate decarboxylase: characterization, investigation of the mevalonate diphosphate binding site, and crystal structure. Voynova NE, et al. Arch Biochem Biophys, 2008 Dec 1. PMID 18823933. Multiple genetic variants along candidate pathways influence plasma high-density lipoprotein cholesterol concentrations. Lu Y, et al. J Lipid Res, 2008 Dec. PMID 18660489. A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. Cell, 2005 Sep 23. PMID 16169070.