

**NFIC Antibody (monoclonal) (M01)**

Mouse monoclonal antibody raised against a full length recombinant NFIC.

Catalog # AT3033a

**Specification**

---

**NFIC Antibody (monoclonal) (M01) - Product Information**

Application	IF, WB, IHC
Primary Accession	<a href="#">P08651</a>
Other Accession	<a href="#">BC012120</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	55675

**NFIC Antibody (monoclonal) (M01) - Additional Information**

Gene ID 4782

**Other Names**

Nuclear factor 1 C-type, NF1-C, Nuclear factor 1/C, CCAAT-box-binding transcription factor, CTF, Nuclear factor I/C, NF-I/C, NFI-C, TGGCA-binding protein, NFIC, NFI

**Target/Specificity**

NFIC (AAH12120, 1 a.a. ~ 428 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

NFIC Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**NFIC Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### NFIC Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to NFIC on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (72.82 KDa) .




NFIC monoclonal antibody (M01), clone 1C12-2A2 Western Blot analysis of NFIC expression in

A-431 ( (Cat # AT3033a )




Western Blot analysis of NFIC expression in transfected 293T cell line by NFIC monoclonal antibody (M01), clone 1C12-2A2.

Lane 1: NFIC transfected lysate(47.6 KDa).  
Lane 2: Non-transfected lysate.



Western blot analysis of NFIC over-expressed 293 cell line, cotransfected with NFIC Validated Chimera RNAi ( (Cat # AT3033a )



Immunoperoxidase of monoclonal antibody to NFIC on formalin-fixed paraffin-embedded human salivary gland. [antibody concentration 3 ug/ml]

#### **NFIC Antibody (monoclonal) (M01) - References**

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Identification of a region in the human IRS2 promoter essential for stress induced transcription depending on SP1, NFI binding and ERK activation in HepG2 cells. Udelhoven M, et al. J Mol Endocrinol, 2010 Feb. PMID 19755487. Nuclear factor I-C links platelet-derived growth factor and transforming growth factor beta1 signaling to skin wound healing progression. Plasari G, et al. Mol Cell Biol, 2009 Nov. PMID 19752192. Regulation of the CYP3A4 and CYP3A7 promoters by members of the nuclear factor I transcription factor family. Riffel AK, et al. Mol Pharmacol, 2009 Nov. PMID 19706729. Nuclear factor I transcription factors regulate IGF binding protein 5 gene transcription in human osteoblasts. Perez-Casellas LA, et al. Biochim Biophys Acta, 2009 Feb. PMID 18809517.