

## NR4A2 Antibody (monoclonal) (M07)

Mouse monoclonal antibody raised against a partial recombinant NR4A2.

Catalog # AT3107a

### Specification

---

#### NR4A2 Antibody (monoclonal) (M07) - Product Information

Application	IF, WB, E
Primary Accession	<a href="#">P43354</a>
Other Accession	<a href="#">BC066890</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	66591

#### NR4A2 Antibody (monoclonal) (M07) - Additional Information

Gene ID 4929

##### Other Names

Nuclear receptor subfamily 4 group A member 2, Immediate-early response protein NOT, Orphan nuclear receptor NURR1, Transcriptionally-inducible nuclear receptor, NR4A2, NOT, NURR1, TINUR

##### Target/Specificity

NR4A2 (AAH66890, 71 a.a. ~ 170 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

##### Dilution

WB~~1:500~1000

##### Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

##### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

##### Precautions

NR4A2 Antibody (monoclonal) (M07) is for research use only and not for use in diagnostic or therapeutic procedures.


#### NR4A2 Antibody (monoclonal) (M07) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### NR4A2 Antibody (monoclonal) (M07) - Images



Immunofluorescence of monoclonal antibody to NR4A2 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



Western Blot analysis of NR4A2 expression in transfected 293T cell line by NR4A2 monoclonal antibody (M07), clone 4A6.

Lane 1: NR4A2 transfected lysate (Predicted MW: 66.6 KDa).  
Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged NR4A2 is approximately 3ng/ml as a capture antibody.

### **NR4A2 Antibody (monoclonal) (M07) - Background**

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcription factor. Mutations in this gene have been associated with disorders related to dopaminergic dysfunction, including Parkinson disease, schizophrenia, and manic depression. Misregulation of this gene may be associated with rheumatoid arthritis. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

### **NR4A2 Antibody (monoclonal) (M07) - References**

1. Neural differentiation potential of human bone marrow-derived mesenchymal stromal cells: misleading marker gene expression. Montzka K, Lassonczyk N, Tschoke B, Neuss S, Fuhrmann T, Franzen R, Smeets R, Brook GA, Woltje M. BMC Neurosci. 2009 Mar 3;10:16.