

PCNA Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a full length recombinant PCNA.

Catalog # AT3236a

Specification

PCNA Antibody (monoclonal) (M02) - Product Information

Application	IF, IP, WB, E
Primary Accession	P12004
Other Accession	BC000491
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a kappa
Calculated MW	28769

PCNA Antibody (monoclonal) (M02) - Additional Information**Gene ID** 5111**Other Names**

Proliferating cell nuclear antigen, PCNA, Cyclin, PCNA

Target/Specificity

PCNA (AAH00491, 1 a.a. ~ 261 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PCNA Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


PCNA Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PCNA Antibody (monoclonal) (M02) - Images



Immunofluorescence of monoclonal antibody to PCNA on HeLa cell. [antibody concentration 10 ug/ml]




Immunoprecipitation of PCNA transfected lysate using anti-PCNA monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with PCNA MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen

(54.45 KDa) .




PCNA monoclonal antibody (M02), clone 1G7 Western Blot analysis of PCNA expression in HeLa (Cat # AT3236a)



Western Blot analysis of PCNA expression in transfected 293T cell line by PCNA monoclonal antibody (M02), clone 1G7.

Lane 1: PCNA transfected lysate(28.8 KDa).
 Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged PCNA is approximately 0.1ng/ml as a capture antibody.

PCNA Antibody (monoclonal) (M02) - Background

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase

delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

PCNA Antibody (monoclonal) (M02) - References

1. The RAD51-stimulatory compound RS-1 can exploit the RAD51 overexpression that exists in cancer cells and tumors. Mason JM, Logan HL, Budke B, Wu M, Pawlowski M, Weichselbaum RR, Kozikowski AP, Bishop DK, Connell PP. *Cancer Res.* 2014 Apr 21.