

PNPLA2 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant PNPLA2.****Catalog # AT3363a****Specification**

PNPLA2 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O96AD5
Other Accession	NM_020376
Reactivity	Human, Rat
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	55316

PNPLA2 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 57104**Other Names**

Patatin-like phospholipase domain-containing protein 2, Adipose triglyceride lipase, Calcium-independent phospholipase A2, Desnutrin, IPLA2-zeta, Pigment epithelium-derived factor, TTS22, Transport-secretion protein 2, TTS2, PNPLA2, ATGL

Target/Specificity

PNPLA2 (NP_065109, 347 a.a. ~ 446 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PNPLA2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PNPLA2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PNPLA2 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



PNPLA2 monoclonal antibody (M01), clone 2H1. Western Blot analysis of PNPLA2 expression in PC-12.



Detection limit for recombinant GST tagged PNPLA2 is approximately 3ng/ml as a capture antibody.

PNPLA2 Antibody (monoclonal) (M01) - Background

This gene encodes an enzyme which catalyzes the first step in the hydrolysis of triglycerides in adipose tissue. Mutations in this gene are associated with neutral lipid storage disease with myopathy.

PNPLA2 Antibody (monoclonal) (M01) - References

High frequency of ETFDH c.250G>A mutation in Taiwanese patients with late-onset lipid storage myopathy. Lan MY, et al. Clin Genet, 2010 Mar 29. PMID 20370797. Rare ATGL haplotypes are associated with increased plasma triglyceride concentrations in the Greenland Inuit. Johansen CT, et al. Int J Circumpolar Health, 2010 Feb. PMID 20167152. Chronic TNF α and cAMP pre-treatment of human adipocytes alter HSL, ATGL and perilipin to regulate basal and stimulated lipolysis. Bezaire V, et al. FEBS Lett, 2009 Sep 17. PMID 19695247. Characterization of desnutrin functional domains: critical residues for triacylglycerol hydrolysis in cultured cells. Duncan RE, et al. J Lipid Res, 2010 Feb. PMID 19692632. Contribution of adipose triglyceride lipase and hormone-sensitive lipase to lipolysis in hMADS adipocytes. Bezaire V, et al. J Biol Chem, 2009 Jul 3. PMID 19433586.