

**RELA Antibody (monoclonal) (M01)**

Mouse monoclonal antibody raised against a partial recombinant RELA.

Catalog # AT3616a

**Specification**

---

**RELA Antibody (monoclonal) (M01) - Product Information**

Application	IF, WB, E
Primary Accession	<a href="#">Q04206</a>
Other Accession	<a href="#">NM_021975</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Lambda
Calculated MW	60219

**RELA Antibody (monoclonal) (M01) - Additional Information****Gene ID** 5970**Other Names**

Transcription factor p65, Nuclear factor NF-kappa-B p65 subunit, Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3, RELA, NFKB3

**Target/Specificity**

RELA (NP\_068810, 432 a.a. ~ 505 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

RELA Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**RELA Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### RELA Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to RELA on HeLa cell. [antibody concentration 35 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (33.88 KDa) .




RELA monoclonal antibody (M01), clone 8G3 Western Blot analysis of RELA expression in HeLa S3

NE ( (Cat # AT3616a )



Detection limit for recombinant GST tagged RELA is 0.03 ng/ml as a capture antibody.



Proximity Ligation Analysis of protein-protein interactions between IKBKB and RELA. HeLa cells were stained with anti- $\text{IKBKB}$  rabbit purified polyclonal 1:1200 and anti- $\text{RELA}$  mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

### **RELA Antibody (monoclonal) (M01) - Background**

$\text{NFKB1}$  (MIM 164011) or  $\text{NFKB2}$  (MIM 164012) is bound to  $\text{REL}$  (MIM 164910),  $\text{RELA}$ , or  $\text{RELB}$  (MIM 604758) to form the  $\text{NFKB}$  complex. The  $\text{p50}$  ( $\text{NFKB1}$ )/ $\text{p65}$  ( $\text{RELA}$ ) heterodimer is the most abundant form of  $\text{NFKB}$ . The  $\text{NFKB}$  complex is inhibited by I-kappa-B proteins ( $\text{NFKBIA}$ , MIM 164008 or  $\text{NFKBIB}$ , MIM 604495), which inactivate  $\text{NFKB}$  by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases ( $\text{IKBKA}$ , MIM 600664, or  $\text{IKBKB}$ , MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the  $\text{NFKB}$  complex. Activated  $\text{NFKB}$  complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).

### **RELA Antibody (monoclonal) (M01) - References**

Variation at the  $\text{NFATC2}$  Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. *Diabetes Care*, 2010 Jul 13. PMID 20628086. Involvement of the  $\text{p65/RelA}$  subunit of  $\text{NF-kappaB}$  in  $\text{TNF-alpha}$ -induced  $\text{SIRT1}$  expression in vascular smooth muscle cells. Zhang HN, et al. *Biochem Biophys Res Commun*, 2010 Jul 2. PMID 20617556. Dichotomy in  $\text{NF-kappaB}$  signaling and chemoresistance in immunoglobulin variable heavy-chain-mutated versus unmutated CLL cells upon  $\text{CD40/TLR9}$  triggering. Tromp JM, et al. *Oncogene*, 2010 Sep 9. PMID 20581863. Nuclear factor kappaB transcription factors are coexpressed and convey a poor outcome in ovarian cancer. Annunziata CM, et al. *Cancer*, 2010 Jul 1. PMID 20564628. Inhibition of  $\text{NFkappaB}$  and pancreatic cancer cell and tumor growth by curcumin is dependent on specificity protein down-regulation.

Jutooru I, et al. J Biol Chem, 2010 Aug 13. PMID 20538607.