

SART2 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant DSE.

Catalog # AT3773a

Specification

SART2 Antibody (monoclonal) (M03) - Product Information

Application	WB, E
Primary Accession	O9UL01
Other Accession	NM_013352
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	109773

SART2 Antibody (monoclonal) (M03) - Additional Information

Gene ID 29940

Other Names

Dermatan-sulfate epimerase, DS epimerase, Chondroitin-glucuronate 5-epimerase, Squamous cell carcinoma antigen recognized by T-cells 2, SART-2, DSE, SART2

Target/Specificity

DSE (NP_037484, 574 a.a. ~ 673 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

SART2 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


SART2 Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SART2 Antibody (monoclonal) (M03) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Western Blot analysis of DSE expression in transfected 293T cell line by DSE monoclonal antibody (M03), clone 6D4.

Lane 1: DSE transfected lysate (Predicted MW: 109.8 KDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged DSE is approximately 0.3ng/ml as a capture antibody.

SART2 Antibody (monoclonal) (M03) - Background

The protein encoded by this gene is a tumor-rejection antigen. This antigen possesses tumor epitopes capable of inducing HLA-A24-restricted and tumor-specific cytotoxic T lymphocytes in cancer patients and may be useful for specific immunotherapy. This gene product is localized to the endoplasmic reticulum. Two transcript variants encoding the same protein have been found for this gene.

SART2 Antibody (monoclonal) (M03) - References

Identification of the active site of DS-epimerase 1 and requirement of N-glycosylation for enzyme function. Pacheco B, et al. J Biol Chem, 2009 Jan 16. PMID 19004833. Biosynthesis of dermatan sulfate: chondroitin-glucuronate C5-epimerase is identical to SART2. Maccarana M, et al. J Biol Chem, 2006 Apr 28. PMID 16505484. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.