

SERPINA6 Antibody (monoclonal) (M04)

Mouse monoclonal antibody raised against a full length recombinant SERPINA6.

Catalog # AT3820a

Specification

SERPINA6 Antibody (monoclonal) (M04) - Product Information

Application	WB, E
Primary Accession	P08185
Other Accession	BC058021
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	45141

SERPINA6 Antibody (monoclonal) (M04) - Additional Information

Gene ID 866

Other Names

Corticosteroid-binding globulin, CBG, Serpin A6, Transcortin, SERPINA6, CBG

Target/Specificity

SERPINA6 (AAH58021, 23 a.a. ~ 405 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

SERPINA6 Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.


SERPINA6 Antibody (monoclonal) (M04) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SERPINA6 Antibody (monoclonal) (M04) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (67.87 KDa) .



Detection limit for recombinant GST tagged SERPINA6 is approximately 1ng/ml as a capture antibody.

SERPINA6 Antibody (monoclonal) (M04) - Background

This gene encodes an alpha-globulin protein with corticosteroid-binding properties. This is the major transport protein for glucocorticoids and progestins in the blood of most vertebrates. The gene localizes to a chromosomal region containing several closely related serine protease inhibitors which may have evolved by duplication events.

SERPINA6 Antibody (monoclonal) (M04) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Genetic variation in neuroendocrine genes associates with somatic symptoms in the general population: results from the EPIFUND study. Holliday KL, et al. J Psychosom Res, 2010 May. PMID 20403506. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Plasma sex hormone-binding globulin, corticosteroid-binding globulin, cortisol, and free cortisol levels in outpatients attending a lipid disorders clinic: a cross-sectional study of 1137 subjects. Lewis JG, et al. Horm Metab Res, 2010 Apr. PMID 20354921. Effect of mutations of the human serpin protein corticosteroid-binding globulin

on cortisol-binding, thermal and protease sensitivity. Braun BC, et al. J Steroid Biochem Mol Biol, 2010 May. PMID 20226861.