

**SIRT1 Antibody (monoclonal) (M03)**

Mouse monoclonal antibody raised against a partial recombinant SIRT1.

Catalog # AT3890a

**Specification**

---

**SIRT1 Antibody (monoclonal) (M03) - Product Information**

Application	WB, E
Primary Accession	<a href="#">O96EB6</a>
Other Accession	<a href="#">BC012499</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	81681

**SIRT1 Antibody (monoclonal) (M03) - Additional Information**

Gene ID 23411

**Other Names**

NAD-dependent protein deacetylase sirtuin-1, hSIRT1, 351-, Regulatory protein SIR2 homolog 1, SIR2-like protein 1, hSIR2, SirtT1 75 kDa fragment, 75SirT1, SIRT1, SIR2L1

**Target/Specificity**

SIRT1 (AAH12499, 456 a.a. ~ 555 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 kDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

SIRT1 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


**SIRT1 Antibody (monoclonal) (M03) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**SIRT1 Antibody (monoclonal) (M03) - Images**



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



SIRT1 monoclonal antibody (M03), clone 7C2. Western Blot analysis of SIRT1 expression in Hela S3 NE ( (Cat # AT3890a )



Detection limit for recombinant GST tagged SIRT1 is 0.1 ng/ml as a capture antibody.

**SIRT1 Antibody (monoclonal) (M03) - Background**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.

### **SIRT1 Antibody (monoclonal) (M03) - References**

SIRT1 regulates UV-induced DNA repair through deacetylating XPA. Fan W, et al. Mol Cell, 2010 Jul 30. PMID 20670893. SIRT1 promotes the central adaptive response to diet restriction through activation of the dorsomedial and lateral nuclei of the hypothalamus. Satoh A, et al. J Neurosci, 2010 Jul 28. PMID 20668205. SIRT1 variants are associated with aging in a healthy Han Chinese population. Zhang WG, et al. Clin Chim Acta, 2010 Nov 11. PMID 20633545. Sirtuin 1 regulation of developmental genes during differentiation of stem cells. Calvanese V, et al. Proc Natl Acad Sci U S A, 2010 Aug 3. PMID 20631301. MicroRNA-34a regulation of endothelial senescence. Ito T, et al. Biochem Biophys Res Commun, 2010 Aug 6. PMID 20627091.