

Tel: 858.875.1900 Fax: 858.875.1999

SMAD1 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a full length recombinant SMAD1. Catalog # AT3937a

Specification

SMAD1 Antibody (monoclonal) (M03) - Product Information

Application **Primary Accession** Other Accession Reactivity

Host Clonality Isotype

Calculated MW

IF, WB, IHC, E 015797

BC001878

Human, Mouse, Rat

mouse Monoclonal IgG1 Kappa 52260

SMAD1 Antibody (monoclonal) (M03) - Additional Information

Gene ID 4086

Other Names

Mothers against decapentaplegic homolog 1, MAD homolog 1, Mothers against DPP homolog 1, JV4-1, Mad-related protein 1, SMAD family member 1, SMAD 1, Smad1, hSMAD1, Transforming growth factor-beta-signaling protein 1, BSP-1, SMAD1, BSP1, MADH1, MADR1

Target/Specificity

SMAD1 (AAH01878.1, 1 a.a. ~ 465 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

SMAD1 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

SMAD1 Antibody (monoclonal) (M03) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides



- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


SMAD1 Antibody (monoclonal) (M03) - Images



Immunofluorescence of monoclonal antibody to SMAD1 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (76.89 KDa) .







SMAD1 monoclonal antibody (M03), clone 2E9 Western Blot analysis of SMAD1 expression in HeLa ((Cat # AT3937a)



SMAD1 monoclonal antibody (M03), clone 2E9. Western Blot analysis of SMAD1 expression in PC-12 ((Cat # AT3937a)




SMAD1 monoclonal antibody (M03), clone 2E9. Western Blot analysis of SMAD1 expression in IMR-32 ((Cat # AT3937a)




SMAD1 monoclonal antibody (M03), clone 2E9. Western Blot analysis of SMAD1 expression in Raw 264.7 ((Cat # AT3937a)






SMAD1 monoclonal antibody (M03), clone 2E9. Western Blot analysis of SMAD1 expression in NIH/3T3 ((Cat # AT3937a)



Immunoperoxidase of monoclonal antibody to SMAD1 on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged SMAD1 is approximately 1ng/ml as a capture antibody.

SMAD1 Antibody (monoclonal) (M03) - Background

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its function in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as SMURF1 and





Tel: 858.875.1900 Fax: 858.875.1999

SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed.

SMAD1 Antibody (monoclonal) (M03) - References

1.Temporal and regional patterns of Smad activation in the rat hippocampus following global ischemia.Nakajima T, Yanagihara M, Nishii HJ Neurol Sci. 2013 Nov 19. pii: S0022-510X(13)03041-4. doi: 10.1016/j.jns.2013.11.012.