

**TAF7 Antibody (monoclonal) (M02)**

Mouse monoclonal antibody raised against a partial recombinant TAF7.

Catalog # AT4148a

**Specification**

---

**TAF7 Antibody (monoclonal) (M02) - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q15545</a>
Other Accession	<a href="#">BC032737</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	40259

**TAF7 Antibody (monoclonal) (M02) - Additional Information**

Gene ID 6879

**Other Names**

Transcription initiation factor TFIID subunit 7, RNA polymerase II TBP-associated factor subunit F, Transcription initiation factor TFIID 55 kDa subunit, TAF(II)55, TAFII-55, TAFII55, TAF7, TAF2F, TAFII55

**Target/Specificity**

TAF7 (AAH32737, 130 a.a. ~ 224 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

TAF7 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


**TAF7 Antibody (monoclonal) (M02) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**TAF7 Antibody (monoclonal) (M02) - Images**



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.08 KDa) .



TAF7 monoclonal antibody (M02), clone 3G6 Western Blot analysis of TAF7 expression in Hela S3 NE ( (Cat # AT4148a )



Detection limit for recombinant GST tagged TAF7 is approximately 0.1ng/ml as a capture antibody.

### **TAF7 Antibody (monoclonal) (M02) - Background**

The intronless gene for this transcription coactivator is located between the protocadherin beta and gamma gene clusters on chromosome 5. The protein encoded by this gene is a component of the TFIID protein complex, a complex which binds to the TATA box in class II promoters and recruits RNA polymerase II and other factors. This particular subunit interacts with the largest TFIID subunit, as well as multiple transcription activators. The protein is required for transcription by promoters targeted by RNA polymerase II.

### **TAF7 Antibody (monoclonal) (M02) - References**

TFIID component TAF7 functionally interacts with both TFIIH and P-TEFb. Gegonne A, et al. Proc Natl Acad Sci U S A, 2008 Apr 8. PMID 18391197. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. Systematic analysis of the protein interaction network for the human transcription machinery reveals the identity of the 7SK capping enzyme. Jeronimo C, et al. Mol Cell, 2007 Jul 20. PMID 17643375. Presence of the anti-leukemic nucleotide analog, 2-chloro-2'-deoxyadenosine-5'-monophosphate, in a promoter sequence alters DNA binding of TATA-binding protein (TBP). Hartman WR, et al. Arch Biochem Biophys, 2007 Mar 15. PMID 17320040. Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983.