

TCF4 Antibody (monoclonal) (M04)

Mouse monoclonal antibody raised against a full length recombinant TCF4.

Catalog # AT4189a

Specification

TCF4 Antibody (monoclonal) (M04) - Product Information

Application	IF, WB
Primary Accession	P15884
Other Accession	BC031056.1
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	71308

TCF4 Antibody (monoclonal) (M04) - Additional Information

Gene ID 6925

Other Names

Transcription factor 4, TCF-4, Class B basic helix-loop-helix protein 19, bHLHb19, Immunoglobulin transcription factor 2, ITF-2, SL3-3 enhancer factor 2, SEF-2, TCF4, BHLHB19, ITF2, SEF2

Target/Specificity

TCF4 (AAH31056.1, 1 a.a. ~ 365 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TCF4 Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.


TCF4 Antibody (monoclonal) (M04) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TCF4 Antibody (monoclonal) (M04) - Images



Immunofluorescence of monoclonal antibody to TCF4 on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (65.89 KDa) .



TCF4 monoclonal antibody (M04), clone 3H4 Western Blot analysis of TCF4 expression in A-549 (

(Cat # AT4189a)

TCF4 Antibody (monoclonal) (M04) - Background

This gene encodes transcription factor 4, a basic helix-turn-helix transcription factor. The encoded protein recognizes an Ephrussi-box ('E-box') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is expressed predominantly in pre-B-cells, although it is found in other tissues as well. Multiple alternatively spliced transcript variants that encode different proteins have been described. [provided by RefSeq]