

## TRIM33 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant TRIM33.

Catalog # AT4352a

### Specification

---

#### TRIM33 Antibody (monoclonal) (M02) - Product Information

Application	IF, WB
Primary Accession	<a href="#">O9UPN9</a>
Other Accession	<a href="#">NM_015906</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	122533

#### TRIM33 Antibody (monoclonal) (M02) - Additional Information

Gene ID 51592

##### Other Names

E3 ubiquitin-protein ligase TRIM33, 632-, Ectodermin homolog, RET-fused gene 7 protein, Protein Rfg7, Transcription intermediary factor 1-gamma, TIF1-gamma, Tripartite motif-containing protein 33, TRIM33, KIAA1113, RFG7, TIF1G

##### Target/Specificity

TRIM33 (NP\_056990, 1006 a.a. ~ 1105 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

##### Dilution

WB~~1:500~1000

##### Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

##### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

##### Precautions

TRIM33 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


#### TRIM33 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TRIM33 Antibody (monoclonal) (M02) - Images



Immunofluorescence of monoclonal antibody to TRIM33 on HeLa cell. [antibody concentration 10 ug/ml]



TRIM33 monoclonal antibody (M02), clone 6F4 Western Blot analysis of TRIM33 expression in HeLa S3 NE ( Cat # L013V3 ).

### TRIM33 Antibody (monoclonal) (M02) - Background

The protein encoded by this gene is thought to be a transcriptional corepressor. However, molecules that interact with this protein have not yet been identified. The protein is a member of the tripartite motif family. This motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. Three alternatively spliced transcript variants for this gene have been described, however, the full-length nature of one variant has not been determined.

### TRIM33 Antibody (monoclonal) (M02) - References

TIF1gamma controls erythroid cell fate by regulating transcription elongation. Bai X, et al. Cell, 2010 Jul 9. PMID 20603019. Transcription intermediary factor 1gamma decreases protein expression of the transcriptional cofactor, LIM-domain-binding 1. Howard PW, et al. Biochem Biophys Res Commun, 2010 Jun 4. PMID 20447379. Inactivation of TIF1gamma cooperates with Kras to induce

cystic tumors of the pancreas. Vincent DF, et al. PLoS Genet, 2009 Jul. PMID 19629168. FAM/USP9x, a deubiquitinating enzyme essential for TGFbeta signaling, controls Smad4 monoubiquitination. Dupont S, et al. Cell, 2009 Jan 9. PMID 19135894. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.