

**TYK2 Antibody (monoclonal) (M02)**

Mouse monoclonal antibody raised against a partial recombinant TYK2.

Catalog # AT4424a

**Specification**

---

**TYK2 Antibody (monoclonal) (M02) - Product Information**

Application	IF, WB, IHC
Primary Accession	<a href="#">P29597</a>
Other Accession	<a href="#">BC014243</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	133650

**TYK2 Antibody (monoclonal) (M02) - Additional Information****Gene ID** 7297**Other Names**

Non-receptor tyrosine-protein kinase TYK2, TYK2

**Target/Specificity**

TYK2 (AAH14243, 276 a.a. ~ 375 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

TYK2 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


**TYK2 Antibody (monoclonal) (M02) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### TYK2 Antibody (monoclonal) (M02) - Images



Immunofluorescence of monoclonal antibody to TYK2 on HeLa cell. [antibody concentration 10 ug/ml]



TYK2 monoclonal antibody (M02), clone 5A4 Western Blot analysis of TYK2 expression in HeLa S3 NE ( Cat # L013V3 ).



Immunoperoxidase of monoclonal antibody to TYK2 on formalin-fixed paraffin-embedded human leiomyosarcoma. [antibody concentration 3 ug/ml]

### TYK2 Antibody (monoclonal) (M02) - Background

This gene encodes a member of the tyrosine kinase and, more specifically, the Janus kinases (JAKs) protein families. This protein associates with the cytoplasmic domain of type I and type II cytokine receptors and promulgate cytokine signals by phosphorylating receptor subunits. It is also component of both the type I and type III interferon signaling pathways. As such, it may play a role in anti-viral immunity. A mutation in this gene has been associated with hyperimmunoglobulin E syndrome (HIES) - a primary immunodeficiency characterized by elevated serum immunoglobulin E.

#### **TYK2 Antibody (monoclonal) (M02) - References**

Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Dengue hemorrhagic fever is associated with polymorphisms in JAK1. Silva LK, et al. Eur J Hum Genet, 2010 Jun 30. PMID 20588308. Interleukin-9 polymorphism in infants with respiratory syncytial virus infection: an opposite effect in boys and girls. Schuurhof A, et al. Pediatr Pulmonol, 2010 Jun. PMID 20503287. Structural and thermodynamic characterization of the TYK2 and JAK3 kinase domains in complex with CP-690550 and CMP-6. Chrencik JE, et al. J Mol Biol, 2010 Jul 16. PMID 20478313. Polymorphisms in innate immunity genes and risk of childhood leukemia. Han S, et al. Hum Immunol, 2010 Jul. PMID 20438785.