

**ULK2 Antibody (monoclonal) (M10)**

Mouse monoclonal antibody raised against a partial recombinant ULK2.

Catalog # AT4465a

**Specification**

---

**ULK2 Antibody (monoclonal) (M10) - Product Information**

|                   |                          |
|-------------------|--------------------------|
| Application       | WB, E                    |
| Primary Accession | <a href="#">Q8IYT8</a>   |
| Other Accession   | <a href="#">BC034988</a> |
| Reactivity        | Human                    |
| Host              | mouse                    |
| Clonality         | Monoclonal               |
| Isotype           | IgG2a Kappa              |
| Calculated MW     | 112694                   |

**ULK2 Antibody (monoclonal) (M10) - Additional Information**

Gene ID 9706

**Other Names**

Serine/threonine-protein kinase ULK2, Unc-51-like kinase 2, ULK2, KIAA0623

**Target/Specificity**

ULK2 (AAH34988, 743 a.a. ~ 843 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

ULK2 Antibody (monoclonal) (M10) is for research use only and not for use in diagnostic or therapeutic procedures.


**ULK2 Antibody (monoclonal) (M10) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ULK2 Antibody (monoclonal) (M10) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.52 KDa) .



Detection limit for recombinant GST tagged ULK2 is approximately 0.03ng/ml as a capture antibody.

### ULK2 Antibody (monoclonal) (M10) - Background

This gene encodes a protein that is similar to a serine/threonine kinase in *C. elegans* which is involved in axonal elongation. The structure of this protein is similar to the *C. elegans* protein in that both proteins have an N-terminal kinase domain, a central proline/serine rich (PS) domain, and a C-terminal (C) domain. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Alternatively spliced transcript variants encoding the same protein have been identified.

### ULK2 Antibody (monoclonal) (M10) - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. *Mol Med*, 2010 Jul-Aug. PMID 20379614. ULK-Atg13-FIP200 complexes mediate mTOR signaling to the autophagy machinery. Jung CH, et al. *Mol Biol Cell*, 2009 Apr. PMID 19225151. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. *Mol Cell Proteomics*, 2008 Mar. PMID 18029348. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. *Genome Res*, 2006 Jan. PMID 16344560. A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. *Cell*, 2005 Sep 23. PMID 16169070.