

USF1 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant USF1.

Catalog # AT4477a

Specification

USF1 Antibody (monoclonal) (M02) - Product Information

Application	WB, E
Primary Accession	P22415
Other Accession	NM_007122
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	33538

USF1 Antibody (monoclonal) (M02) - Additional Information

Gene ID 7391

Other Names

Upstream stimulatory factor 1, Class B basic helix-loop-helix protein 11, bHLHb11, Major late transcription factor 1, USF1, BHLHB11, USF

Target/Specificity

USF1 (NP_009053, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

USF1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


USF1 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


USF1 Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



USF1 monoclonal antibody (M02), clone 2A7 Western Blot analysis of USF1 expression in MCF-7 (Cat # AT4477a)



Detection limit for recombinant GST tagged USF1 is approximately 0.1ng/ml as a capture antibody.

USF1 Antibody (monoclonal) (M02) - Background

This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Two transcript variants encoding distinct isoforms have been identified for this gene.

USF1 Antibody (monoclonal) (M02) - References

1.Association of thymidylate synthase enhancer region polymorphisms with thymidylate synthase activity in vivo.de Bock CE, Garg MB, Scott N, Sakoff JA, Scorgie FE, Ackland SP, Lincz LF.Pharmacogenomics J. 2010 Jun 8. [Epub ahead of print]2.Use of a single chain antibody library for ovarian cancer biomarker discovery.Ramirez AB, Loch CM, Zhang Y, Liu Y, Wang X, Wayner EA, Sargent JE, Sibani S, Hainsworth E, Mendoza EA, Eugene R, Labaer J, Urban ND, McIntosh MW, Lampe PD.Mol Cell Proteomics. 2010 May 13. [Epub ahead of print]3.A role of DNA-PK for the metabolic gene regulation in response to insulin.Wong RH, Chang I, Hudak CS, Hyun S, Kwan HY, Sul HS.Cell. 2009 Mar 20;136(6):1056-72.