

APC Anti-Human CD45 (HI30) Antibody

Catalog # ATB10029

Specification

APC Anti-Human CD45 (HI30) Antibody - Product Information

Application FC

Isotype Mouse IgG1, kappa
Concentration 5 uL (0.125 ug)/test

Reactivity Human

Formulation 10 mM NaH2PO4, 150 mM NaCl, 0.09%

NaN3, 0.1% gelatin, pH7.2

Host Mouse

APC Anti-Human CD45 (HI30) Antibody - Additional Information

Gene ID 5788
Gene Name PTPRC

Alternative Name(s)

Leukocyte Common Antigen, LCA, Ly-5

Format

APC

Preparation

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Application Notes

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 uL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 uL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10e5 to 1x10e8 cells.

Storage Conditions

2-8°C protected from light

APC Anti-Human CD45 (HI30) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence




- Immunoprecipitation
- Flow Cytomety
- Cell Culture

APC Anti-Human CD45 (HI30) Antibody - Images



Human peripheral blood lymphocytes were stained with 5 uL (0.125 ug) Anti-Human CD45 APC (ATB10029) (solid line) or 0.125 ug Mouse IgG1 APC isotype control.



Human peripheral blood lymphocytes were stained with 5 uL (0.125 ug) APC Anti-Human CD45 (HI30) manufactured by Tonbo Biosciences (left panel) or eBioscience (right panel).

APC Anti-Human CD45 (HI30) Antibody - Background

The HI30 antibody reacts with human CD45, one of the most abundant hematopoietic markers and one that is expressed on all leukocytes (the Leukocyte Common Antigen, LCA). CD45 is a protein tyrosine phosphatase existing in several isoforms, each being generated and expressed in cell-specific patterns. With its broad cell distribution, CD45 is critical for many leukocyte functions, regulating signal transduction and cell activation associated with the T cell receptor, B cell receptor, and IL-2 receptor. Other forms of CD45, with restricted cellular expression, include CD45R (B220), CD45RA, CD45RB, CD45RO and others. The HI30 antibody is widely used as a marker for human CD45 expression on T cells, B cells, monocytes, macrophages, and NK cells.