

**FITC Anti-Human GARP (GARP5) Antibody**  
Catalog # ATB10136

**Specification**

---

**FITC Anti-Human GARP (GARP5) Antibody - Product Information**

|               |   |
|---------------|---|
| Application   | FC  |
| Isotype       | Mouse IgG1  |
| Concentration | 5 uL (0.5 ug)/test  |
| Reactivity    | Human   |
| Formulation   | 10 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, 0.09% Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , 0.1% gelatin, pH7.2 |
| Host          | Mouse   |

**FITC Anti-Human GARP (GARP5) Antibody - Additional Information**

|                     |       |
|---------------------|-------|
| Gene ID             | 2615  |
| Gene Name           | LRR32 |
| Alternative Name(s) |       |
| LRR32, Garpin       |       |

**Format**  
FITC

**Preparation**

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

**Application Notes**

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 uL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 uL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10<sup>5</sup> to 1x10<sup>8</sup> cells.

**Storage Conditions**

2-8°C protected from light


**FITC Anti-Human GARP (GARP5) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **FITC Anti-Human GARP (GARP5) Antibody - Images**



Untransfected (left) or GARP transfected (right) cells were stained with 5 uL (0.5 ug) FITC Anti-Human GARP (ATB10136) (solid line) or 0.5 ug FITC Mouse IgG1 isotype control (dashed line).

### **FITC Anti-Human GARP (GARP5) Antibody - Background**

The GARP5 antibody reacts with human GARP, also known as LRRC32 or Garpin, an 80 kDa transmembrane protein which acts as a receptor for the latent form of TGF-beta 1 (pro-TGF-beta), preventing its secretion. Specifically, GARP is reported to associate with the pro-domain of TGF-beta 1, known as latency-associated peptide (LAP), which must be cleaved to release the biologically active cytokine. The putative role of GARP in sequestering the TGF-beta is important for regulating its activity, as TGF-beta signaling is involved in development and function of Th17, Treg and many other immune cell types. GARP expression has been shown on platelets and at high levels on Foxp3+ Treg cells, where it is proposed to be a phenotypic identifier for activated human Treg cells. The GARP5 antibody may be used for analysis of human GARP by flow cytometry, immunoprecipitation and Western blotting. Please choose the appropriate format for each application.