

**PE-Cy7 Anti-Human CD19 (HIB19) Antibody**  
Catalog # ATB10241**Specification****PE-Cy7 Anti-Human CD19 (HIB19) Antibody - Product Information**

Application	FC
Isotype	Mouse IgG1, kappa
Concentration	5 uL (0.25 ug)/test
Reactivity	Human
Formulation	10 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, 0.09% Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , 0.1% gelatin, pH7.2
Host	Mouse

**PE-Cy7 Anti-Human CD19 (HIB19) Antibody - Additional Information**

Gene ID	930
Gene Name	CD19
Alternative Name(s)	
Leu-12, B4	

**Format**  
PE-Cy7**Preparation**

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

**Application Notes**

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 uL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 uL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10<sup>5</sup> to 1x10<sup>8</sup> cells.

**Storage Conditions**

2-8°C protected from light


**PE-Cy7 Anti-Human CD19 (HIB19) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PE-Cy7 Anti-Human CD19 (HIB19) Antibody - Images



Human peripheral blood lymphocytes were stained with 5  $\mu$ L (0.25  $\mu$ g) PE-Cy7 Anti-Human CD19 (ATB10241) (solid line) or 0.25  $\mu$ g PE-Cy7 Mouse IgG1 isotype control (dashed line).

### PE-Cy7 Anti-Human CD19 (HIB19) Antibody - Background

The HIB19 antibody reacts with human CD19, 95 kDa glycoprotein which acts as a co-receptor, along with CD21, CD81 and CD225, in support of the functional B cell receptor (BCR). This complex provides antigen-specific recognition and subsequent activation of B cells to proliferate and differentiate into antibody-secreting cells (plasma cells) or memory B cells, which are crucial for secondary antigen encounter. CD19 is a lineage-differentiation marker, as its expression is detectable at the earliest B cell stages, through development, and is finally lost upon transition to mature plasma cells. The HIB19 antibody is widely used as a phenotypic marker for CD19 expression on B cells, as well as on dendritic cell subsets.