

HSP90 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5029

Specification

HSP90 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P07900
Other Accession	P02828 , P34058 , P11499 , Q4R4T5 , P08238 , Q04619 , Q76LV1 , P82995 , P30946 , O02705 , P07901 , Q4R4P1 , P46633 , P11501 , Q76LV2 , Q58FF6 , Q58FF7 , Q58FF8 , Q90474 , Q9GKX7 , Q9GKX8 , P30947
Reactivity	Human, Mouse, Rat
Predicted	Zebrafish, Bovine, Chicken, Hamster, Horse, Monkey, Pig, Rabbit, Drosophila
Host	Rabbit
Clonality	polyclonal
Calculated MW	H=85;M=85;Rat=85 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

HSP90 Antibody (N-term) - Additional Information

Gene ID 3320

Antigen Region
57-89

Other Names

Heat shock protein HSP 90-alpha, Heat shock 86 kDa, HSP 86, HSP86, Renal carcinoma antigen NY-REN-38, HSP90AA1, HSP90A, HSPC1, HSPCA

Dilution

WB~~1:1000

Target/Specificity

This HSP90 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 57-89 amino acids from the N-terminal region of human HSP90.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSP90 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic

procedures.

HSP90 Antibody (N-term) - Protein Information

Name HSP90AA1 ([HGNC:5253](#))

Synonyms HSP90A, HSPC1, HSPCA

Function

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity which is essential for its chaperone activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function (PubMed:11274138, PubMed:12526792, PubMed:15577939, PubMed:15937123, PubMed:27353360, PubMed:29127155). Engages with a range of client protein classes via its interaction with various co-chaperone proteins or complexes, that act as adapters, simultaneously able to interact with the specific client and the central chaperone itself (PubMed:29127155). Recruitment of ATP and co-chaperone followed by client protein forms a functional chaperone. After the completion of the chaperoning process, properly folded client protein and co- chaperone leave HSP90 in an ADP-bound partially open conformation and finally, ADP is released from HSP90 which acquires an open conformation for the next cycle (PubMed:26991466, PubMed:27295069). Plays a critical role in mitochondrial import, delivers preproteins to the mitochondrial import receptor TOMM70 (PubMed:12526792). Apart from its chaperone activity, it also plays a role in the regulation of the transcription machinery. HSP90 and its co-chaperones modulate transcription at least at three different levels (PubMed:25973397). In the first place, they alter the steady-state levels of certain transcription factors in response to various physiological cues (PubMed:25973397). Second, they modulate the activity of certain epigenetic modifiers, such as histone deacetylases or DNA methyl transferases, and thereby respond to the change in the environment (PubMed:25973397). Third, they participate in the eviction of histones from the promoter region of certain genes and thereby turn on gene expression (PubMed:25973397). Binds bacterial lipopolysaccharide (LPS) and mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed:11276205). Antagonizes STUB1-mediated inhibition of TGF-beta signaling via inhibition of STUB1-mediated SMAD3 ubiquitination and degradation (PubMed:24613385). Mediates the association of TOMM70 with IRF3 or TBK1 in mitochondrial outer membrane which promotes host antiviral response (PubMed:20628368, PubMed:25609812).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P07901}. Cytoplasm {ECO:0000250|UniProtKB:P07901}.


Melanosome. Cell membrane. Mitochondrion. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

HSP90 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HSP90 Antibody (N-term) - Images



Western blot analysis of lysates from HeLa, A431, A549, mouse NIH/3T3, rat PC-12, SK-N-MC cell line (from left to right), using HSP90 Antibody (N-term)(Cat. #AW5029). AW5029 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

HSP90 Antibody (N-term) - Background

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function.

HSP90 Antibody (N-term) - References

Soeda E., et al. Nucleic Acids Res. 17:7108-7108(1989).
Yamazaki M., et al. Agric. Biol. Chem. 54:3163-3170(1990).
Hickey E., et al. Mol. Cell. Biol. 9:2615-2626(1989).

Chen B., et al. Genomics 86:627-637(2005).
Ota T., et al. Nat. Genet. 36:40-45(2004).