

TAF7 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5093

Specification

TAF7 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O15545
Other Accession	O9R1C0 , O4R5A5 , O6R1L1 , O2HJG8 , NP_005633.2
Reactivity	Human
Predicted	Bovine, Hamster, Monkey, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=40;M=39 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

TAF7 Antibody (C-term) - Additional Information

Gene ID 6879

Antigen Region
308-337

Other Names
TAF7; TAF2F; TAFII55; Transcription initiation factor TFIID subunit 7; RNA polymerase II TBP-associated factor subunit F; Transcription initiation factor TFIID 55 kDa subunit

Dilution
WB~~1:1000

Target/Specificity
This TAF7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 308-337 amino acids from the C-terminal region of human TAF7.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
TAF7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TAF7 Antibody (C-term) - Protein Information

Name TAF7**Synonyms** TAF2F, TAFII55**Function**

The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed:33795473). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:33795473). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:10438527, PubMed:33795473). TAF7 forms a promoter DNA binding subcomplex of TFIID, together with TAF1 and TAF2 (PubMed:33795473). Part of a TFIID complex containing TAF10 (TFIID alpha) and a TFIID complex lacking TAF10 (TFIID beta) (PubMed:10438527).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9R1C0}.


Tissue Location


Ubiquitous.

TAF7 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAF7 Antibody (C-term) - Images



Western blot analysis of lysates from A549, HeLa cell line (from left to right), using TAF7 Antibody (C-term) (Cat. #AW5093). AW5093 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.

TAF7 Antibody (C-term) - Background

The intronless gene for this transcription coactivator is located between the protocadherin beta and gamma gene clusters on chromosome 5. The protein encoded by this gene is a component of the TFIID protein complex, a complex which binds to the TATA box in class II promoters and recruits RNA polymerase II and other factors. This particular subunit interacts with the largest TFIID subunit, as well as multiple transcription activators. The protein is required for transcription by promoters targeted by RNA polymerase II.

TAF7 Antibody (C-term) - References

- Gegonne, A., et al. Proc. Natl. Acad. Sci. U.S.A. 105(14):5367-5372(2008)
- Hartman, W.R., et al. Arch. Biochem. Biophys. 459(2):223-232(2007)
- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Couture, J.F., et al. Nat. Struct. Mol. Biol. 13(2):140-146(2006)