

CRTAP Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5097**Specification**

CRTAP Antibody (N-term) - Product Information

Application	IF, WB,E
Primary Accession	O75718
Other Accession	NP_006362.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=47;M=46 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

CRTAP Antibody (N-term) - Additional Information**Gene ID** 10491**Antigen Region**

31-59

Other Names

CRTAP; CASP; Cartilage-associated protein

Dilution

IF~~1:25

WB~~1:1000

Target/Specificity

This CRTAP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 31-59 amino acids from the N-terminal region of human CRTAP.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CRTAP Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CRTAP Antibody (N-term) - Protein Information

Name CRTAP

Synonyms CASP

Function

Necessary for efficient 3-hydroxylation of fibrillar collagen prolyl residues.

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location


Found in articular chondrocytes. Expressed in a variety of tissues

CRTAP Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRTAP Antibody (N-term) - Images



Fluorescent image of HeLa cells stained with CRTAP Antibody (N-term)(Cat#AW5097). AW5097 was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western blot analysis of lysates from HT1080,ZR-75-1 cell line (from left to right), using CRTAP Antibody (N-term)(Cat. #AW5097). AW5097 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

CRTAP Antibody (N-term) - Background

The protein encoded by this gene is similar to the chicken and mouse CRTAP genes. The encoded protein is a scaffolding protein that may influence the activity of at least one member of the cytohesin/ARNO family in response to specific cellular stimuli. Defects in this gene are associated with osteogenesis imperfecta, a connective tissue disorder characterized by bone fragility and low bone mass.

CRTAP Antibody (N-term) - References

Morello, R., et al. *Curr Osteoporos Rep* 8(2):77-83(2010)
Li, G.H., et al. *Osteoporos Int* 21(6):1009-1020(2010)
Fratzl-Zelman, N., et al. *Bone* 46(3):820-826(2010)
Chang, W., et al. *Hum. Mol. Genet.* 19(2):223-234(2010)
Marini, J.C., et al. *Cell Tissue Res.* 339(1):59-70(2010)