

Phospho-HIST1H3B3(S10) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5153

Specification

Phospho-HIST1H3B3(S10) Antibody - Product Information

Application	WB,E
Primary Accession	P68431
Reactivity	Human
Predicted	Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=15,M=15,Rat=15 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

Phospho-HIST1H3B3(S10) Antibody - Additional Information

Gene ID 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968

Antigen Region

8-35

Other Names

HIST1H3A; H3FA; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l

Dilution

WB~~1:500

Target/Specificity

This HIST1H3B3 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S10 of human HIST1H3B3.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-HIST1H3B3(S10) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-HIST1H3B3(S10) Antibody - Protein Information

Name H3C1 ([HGNC:4766](#))

Synonyms H3FA, HIST1H3A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

Phospho-HIST1H3B3(S10) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-HIST1H3B3(S10) Antibody - Images



Western blot analysis of lysate from HeLa cell line, using Phospho-H3-pS10 Antibody (Cat. #AW5153). AW5153 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.



Western blot analysis of lysate from HeLa cell line, using Phospho-H3-pS10 Antibody (Cat. #AW5153). AW5153 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Phospho-HIST1H3B3(S10) Antibody - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Phospho-HIST1H3B3(S10) Antibody - References

- Lusic, M., et al., EMBO J. 22(24):6550-6561 (2003).
- Deng, L., et al., Virology 289(2):312-326 (2001).
- Deng, L., et al., Virology 277(2):278-295 (2000).
- El Kharroubi, A., et al., Mol. Cell. Biol. 18(5):2535-2544 (1998).
- Albig, W., et al., Hum. Genet. 101(3):284-294 (1997).