

**Phospho-ERBB2(Y1139) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5159****Specification**

---

**Phospho-ERBB2(Y1139) Antibody - Product Information**

Application	WB,E
Primary Accession	<a href="#">P04626</a>
Other Accession	<a href="#">NP_001005862.1</a>
Reactivity	Human
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=138 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**Phospho-ERBB2(Y1139) Antibody - Additional Information****Gene ID** 2064**Antigen Region**  
1114-1146**Other Names**

ERBB2; HER2; MLN19; NEU; NGL; Receptor tyrosine-protein kinase erbB-2; Metastatic lymph node gene 19 protein; Proto-oncogene Neu; Proto-oncogene c-ErbB-2; Tyrosine kinase-type cell surface receptor HER2; p185erbB2; CD\_antigen=CD340; Flags: Precursor

**Dilution**

WB~~1:4000

**Target/Specificity**

This ERBB2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1139 of human ERBB2.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-ERBB2(Y1139) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-ERBB2(Y1139) Antibody - Protein Information**

**Name** ERBB2

**Synonyms** HER2, MLN19, NEU, NGL

**Function**

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Note=Internalized from the cell membrane in response to EGF stimulation. [Isoform 2]: Cytoplasm. Nucleus.

**Tissue Location**


Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.


**Phospho-ERBB2(Y1139) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Phospho-ERBB2(Y1139) Antibody - Images**





Western blot analysis of lysate from A431 cell line, using Phospho-ERBB2-Y1139 Antibody (Cat. #AW5159). AW5159 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.



Western blot analysis of extracts from A431 cells, untreated or treated with EGF, 100ng/ml, using Phospho-ERBB2 Antibody (Y1139) (upper) or Beta-actin (lower).

### Phospho-ERBB2(Y1139) Antibody - Background

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results

in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized.

#### **Phospho-ERBB2(Y1139) Antibody - References**

- Geradts, J., et al. Cancer Invest. 28(9):969-977(2010)  
Zaoui, K., et al. Proc. Natl. Acad. Sci. U.S.A. 107(43):18517-18522(2010)  
Oliveras, G., et al. Ann. N. Y. Acad. Sci. 1210, 86-92 (2010) :  
Han, J.S., et al. Anticancer Res. 30(9):3407-3412(2010)  
Stackiewicz, R., et al. Isr. Med. Assoc. J. 12(5):290-295(2010)