

MGMT Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AW5190

Specification

MGMT Antibody - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P16455
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	H=22 KDa
Isotype	IgG1
Antigen Source	Human

MGMT Antibody - Additional Information

Gene ID 4255

Antigen Region
1-217

Other Names

MGMT;Methylated-DNA--protein-cysteine methyltransferase; Methylated-DNA--protein-cysteine methyltransferase; 6-O-methylguanine-DNA methyltransferase; Methylated-DNA--protein-cysteine methyltransferase; O-6-methylguanine-DNA-alkyltransferase

Dilution

IF~~1:25
WB~~1:1000
IHC-P~~1:25

Target/Specificity

Purified His-tagged MGMT protein was used to produced this monoclonal antibody.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MGMT Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MGMT Antibody - Protein Information

Name MGMT

Function

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) and O4-methylthymine (O4-MeT) in DNA. Repairs the methylated nucleobase in DNA by stoichiometrically transferring the methyl group to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.

Cellular Location


Nucleus.

MGMT Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MGMT Antibody - Images



Fluorescent image of HeLa cells stained with MGMT Antibody (Cat#AW5190). AW5190 was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western blot analysis of lysates from MCF-7, HeLa, Jurkat cell line (from left to right), using MGMT Antibody (Cat. #AW5190). AW5190 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.



Immunohistochemical analysis of paraffin-embedded H. small intestine section using MGMT Antibody (Cat#AW5190). AW5190 was diluted at 1:25 dilution. An undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

MGMT Antibody - Background

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) in DNA. Repairs alkylated guanine in DNA by stoichiometrically transferring the alkyl group at the O-6 position to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.

MGMT Antibody - References

- Edara S., et al. Carcinogenesis 16:1637-1642(1995).
- Tano K., et al. Proc. Natl. Acad. Sci. U.S.A. 87:686-690(1990).
- Rydberg B., et al. J. Biol. Chem. 265:9563-9569(1990).
- Koike G., et al. J. Biol. Chem. 265:14754-14762(1990).
- Hayakawa H., et al. J. Mol. Biol. 213:739-747(1990).