

**SFN Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5213**

**Specification**

---

**SFN Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P31947</a>
Other Accession	<a href="#">O70456</a> , <a href="#">Q0VC36</a> , <a href="#">NP_006133.1</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=28,24;M=28 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**SFN Antibody (Center) - Additional Information**

**Gene ID** 2810

**Antigen Region**  
120-149

**Other Names**  
SFN; HME1; 14-3-3 protein sigma; Epithelial cell marker protein 1; Stratifin

**Dilution**  
WB~~1:1000

**Target/Specificity**  
This SFN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 120-149 amino acids from the Central region of human SFN.

**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**  
SFN Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**SFN Antibody (Center) - Protein Information**

**Name** SFN

**Synonyms** HME1 {ECO:0000303|PubMed:1390337}

### Function

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/15731107" target="\_blank">15731107</a>, PubMed:<a href="http://www.uniprot.org/citations/22634725" target="\_blank">22634725</a>, PubMed:<a href="http://www.uniprot.org/citations/28202711" target="\_blank">28202711</a>, PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:<a href="http://www.uniprot.org/citations/15731107" target="\_blank">15731107</a>, PubMed:<a href="http://www.uniprot.org/citations/22634725" target="\_blank">22634725</a>, PubMed:<a href="http://www.uniprot.org/citations/28202711" target="\_blank">28202711</a>, PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Binding generally results in the modulation of the activity of the binding partner (PubMed:<a href="http://www.uniprot.org/citations/15731107" target="\_blank">15731107</a>, PubMed:<a href="http://www.uniprot.org/citations/22634725" target="\_blank">22634725</a>, PubMed:<a href="http://www.uniprot.org/citations/28202711" target="\_blank">28202711</a>, PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Promotes cytosolic retention of GBP1 GTPase by binding to phosphorylated GBP1, thereby inhibiting the innate immune response (PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Also acts as a TP53/p53-regulated inhibitor of G2/M progression (PubMed:<a href="http://www.uniprot.org/citations/9659898" target="\_blank">9659898</a>). When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity). May also regulate MDM2 autoubiquitination and degradation and thereby activate p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/18382127" target="\_blank">18382127</a>).

### Cellular Location

Cytoplasm, cytosol. Nucleus {ECO:0000250|UniProtKB:O70456}. Secreted. Note=May be secreted by a non-classical secretory pathway.

### Tissue Location


Present mainly in tissues enriched in stratified squamous keratinizing epithelium.


## SFN Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## SFN Antibody (Center) - Images





Western blot analysis of lysates from HeLa cell line, mouse bladder tissue lysate, A431 cell line (from left to right), using SFN Antibody (Center) (Cat. #AW5213). AW5213 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

#### **SFN Antibody (Center) - Background**

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathway. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity).  
p53-regulated inhibitor of G2/M progression.

#### **SFN Antibody (Center) - References**

- Ren, H.Z., et al. Dig. Dis. Sci. 55(9):2552-2560(2010)
- Liu, Y., et al. Brain Res. 1336, 98-102 (2010) :
- Pei, H.P., et al. Zhonghua Wei Chang Wai Ke Za Zhi 13(3):213-215(2010)
- Syrjanen, S., et al. Am. J. Clin. Pathol. 133(2):232-240(2010)
- Zurita, M., et al. BMC Cancer 10, 217 (2010) :