

**CLIC1 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5215**

**Specification**

---

**CLIC1 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O00299</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=27 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**CLIC1 Antibody (Center) - Additional Information**

**Gene ID** 1192

**Antigen Region**  
136-166

**Other Names**  
CLIC1;Chloride intracellular channel protein 1

**Dilution**  
WB~~1:1000

**Target/Specificity**  
This CLIC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 136-166 amino acids from the Central region of human CLIC1.

**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**  
CLIC1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**CLIC1 Antibody (Center) - Protein Information**

**Name** CLIC1

## Synonyms G6, NCC27

### Function

Can insert into membranes and form chloride ion channels. Channel activity depends on the pH. Membrane insertion seems to be redox-regulated and may occur only under oxydizing conditions. Involved in regulation of the cell cycle.

### Cellular Location

Nucleus. Nucleus membrane; Single-pass membrane protein. Cytoplasm. Cell membrane; Single-pass membrane protein. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q6MG61}. Note=Mostly in the nucleus including in the nuclear membrane (PubMed:12681486, PubMed:9139710). Small amount in the cytoplasm and the plasma membrane (PubMed:9139710). Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain (PubMed:11551966, PubMed:11940526, PubMed:12681486, PubMed:14613939, PubMed:9139710). Might not be present in the nucleus of cardiac cells (By similarity) {ECO:0000250|UniProtKB:Q6MG61, ECO:0000269|PubMed:11551966, ECO:0000269|PubMed:11940526, ECO:0000269|PubMed:12681486, ECO:0000269|PubMed:14613939, ECO:0000269|PubMed:9139710}

### Tissue Location


Expression is prominent in heart, placenta, liver, kidney and pancreas.

## CLIC1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CLIC1 Antibody (Center) - Images



Western blot analysis of lysates from human placenta tissue lysate, HL-60, MCF-7, Raji, WiDr cell line (from left to right), using CLIC1 Antibody (Center) (Cat. #AW5215). AW5215 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.

#### **CLIC1 Antibody (Center) - Background**

Can insert into membranes and form chloride ion channels. Channel activity depends on the pH. Membrane insertion seems to be redox-regulated and may occur only under oxidizing conditions. Involved in regulation of the cell cycle.

#### **CLIC1 Antibody (Center) - References**

Xie T., et al. *Genome Res.* 13:2621-2636(2003).  
Shiina S., et al. Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.  
Valenzuela S.M., et al. *J. Biol. Chem.* 272:12575-12582(1997).  
Noh Y.H., et al. Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.  
Chuang J.Z., et al. *J. Neurosci.* 19:2919-2928(1999).