

**Phospho-SHP2(Y584) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5224**

**Specification**

---

**Phospho-SHP2(Y584) Antibody - Product Information**

Application	WB, DB,E
Primary Accession	<a href="#">Q06124</a>
Other Accession	<a href="#">P41499</a> , <a href="#">P35235</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=45 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**Phospho-SHP2(Y584) Antibody - Additional Information**

**Gene ID** 5781

**Antigen Region**  
574-590

**Other Names**  
PTPN11; PTP2C; SHPTP2; Tyrosine-protein phosphatase non-receptor type 11; Protein-tyrosine phosphatase 1D; Protein-tyrosine phosphatase 2C; SH-PTP2; SH-PTP3

**Dilution**  
WB~~1:1000  
DB~~1:500

**Target/Specificity**  
This SHP2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y584 of human SHP2.

**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**  
Phospho-SHP2(Y584) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-SHP2(Y584) Antibody - Protein Information**

**Name** PTPN11

**Synonyms** PTP2C, SHPTP2

### Function

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:<a href="http://www.uniprot.org/citations/10655584" target="\_blank">10655584</a>, PubMed:<a href="http://www.uniprot.org/citations/18559669" target="\_blank">18559669</a>, PubMed:<a href="http://www.uniprot.org/citations/18829466" target="\_blank">18829466</a>, PubMed:<a href="http://www.uniprot.org/citations/26742426" target="\_blank">26742426</a>, PubMed:<a href="http://www.uniprot.org/citations/28074573" target="\_blank">28074573</a>). Positively regulates MAPK signal transduction pathway (PubMed:<a href="http://www.uniprot.org/citations/28074573" target="\_blank">28074573</a>). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:<a href="http://www.uniprot.org/citations/28074573" target="\_blank">28074573</a>). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:<a href="http://www.uniprot.org/citations/18559669" target="\_blank">18559669</a>). Dephosphorylates CDC73 (PubMed:<a href="http://www.uniprot.org/citations/26742426" target="\_blank">26742426</a>). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:<a href="http://www.uniprot.org/citations/19275884" target="\_blank">19275884</a>).

### Cellular Location

Cytoplasm. Nucleus

### Tissue Location


Widely expressed, with highest levels in heart, brain, and skeletal muscle.


### Phospho-SHP2(Y584) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Phospho-SHP2(Y584) Antibody - Images





Western blot analysis of lysates from mouse NIH/3T3 cell line, untreated or treated with PDGF, using Phospho-SHP2-pY584 (Cat. #AW5224) (upper) or Beta-actin (lower).



Dot blot analysis of anti-Phospho-SHP2-pY584 Phospho-specific Pab (Cat. #AW5224) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

### Phospho-SHP2(Y584) Antibody - Background

SHP2 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates. This PTP is widely expressed in most tissues and plays a regulatory role in various cell signaling events that are important for a diversity of cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration.

### Phospho-SHP2(Y584) Antibody - References

- Carver, K.C., et al. J. Biol. Chem. 285(11):8003-8012(2010)
- Pierpont, E.I., et al. Am. J. Med. Genet. A 152A (3), 591-600 (2010)
- Rani, D.S., et al. Mitochondrion 10(2):166-173(2010)
- Bakken, T., et al. Virology 397(2):379-388(2010)