

NRAS Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5296

Specification

NRAS Antibody (C-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P, FC,E |
| Primary Accession | P01111 |
| Other Accession | Q2MJK3 |
| Reactivity | Human |
| Predicted | Pig |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | H=21;M=21;Rat=21 KDa |
| Isotype | Rabbit IgG |
| Antigen Source | HUMAN |

NRAS Antibody (C-term) - Additional Information

Gene ID 4893

Antigen Region
147-179

Other Names
NRAS; HRAS1; GTPase NRas; Transforming protein N-Ras

Dilution
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Target/Specificity
This NRAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 147-179 amino acids from the C-terminal region of human NRAS.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
NRAS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NRAS Antibody (C-term) - Protein Information

Name NRAS

Synonyms HRAS1

Function

Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor
Note=Shuttles between the plasma membrane and the Golgi apparatus

NRAS Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NRAS Antibody (C-term) - Images



Western blot analysis of lysates from MCF-7, A375 cell line (from left to right), using NRAS Antibody (C-term) (Cat. #AW5296). AW5296 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with NRAS antibody (C-term) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



NRAS Antibody (C-term) (Cat. #AW5296) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

NRAS Antibody (C-term) - Background

NRAS is a membrane protein that shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. This protein, which has intrinsic GTPase activity, is activated to a GTP-bound form by a GTPase activating protein and inactivated to a GDP-bound form by a guanine nucleotide-exchange factor. Defects in the gene encoding this protein are a cause of juvenile myelomonocytic leukemia (JMML).

NRAS Antibody (C-term) - References

Smalley,K.S., Cancer Res. 68 (14), 5743-5752 (2008)
Banerji,U., Mol. Cancer Ther. 7 (4), 737-739 (2008)