

**ZNF192 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5302****Specification**

---

**ZNF192 Antibody (Center) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">Q15776</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>H=66 KDa</b>
Isotype	<b>Rabbit IgG</b>
Antigen Source	<b>HUMAN</b>

**ZNF192 Antibody (Center) - Additional Information****Gene ID** 7745**Antigen Region**  
184-206**Other Names**

Zinc finger protein with KRAB and SCAN domains 8, LD5-1, Zinc finger protein 192, ZKSCAN8, ZNF192

**Dilution**

WB~~1:1000

**Target/Specificity**

This ZNF192 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 184-206 amino acids from the Central region of human ZNF192.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ZNF192 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ZNF192 Antibody (Center) - Protein Information****Name** ZKSCAN8

## Synonyms ZNF192

### Function

May be involved in transcriptional regulation.

### Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

## ZNF192 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## ZNF192 Antibody (Center) - Images



Western blot analysis of lysates from HepG2,SH-SY5Y cell line (from left to right), using ZNF192 Antibody (Center)(Cat. #AW5302). AW5302 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

## ZNF192 Antibody (Center) - Background

May be involved in transcriptional regulation.

## ZNF192 Antibody (Center) - References

Lee P.L.,et al.Genomics 43:191-201(1997). Mungall A.J.,et al.Nature 425:805-811(2003). Mural R.J.,et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Beutler E.,et al.Blood Cells Mol. Dis. 21:207-216(1995). Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).