

HMGA2 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5342

Specification

HMGA2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P52926
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=12, 9, 10, 11;M=12 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

HMGA2 Antibody (N-term) - Additional Information

Gene ID 8091

Antigen Region
15-49

Other Names
High mobility group protein HMGI-C, High mobility group AT-hook protein 2, HMGA2, HMGIC

Dilution
WB~~1:1000

Target/Specificity
This HMGA2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 15-49 amino acids from the N-terminal region of human HMGA2.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
HMGA2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HMGA2 Antibody (N-term) - Protein Information

Name HMGA2

Synonyms HMGIC

Function

Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2. Plays an important role in chromosome condensation during the meiotic G2/M transition of spermatocytes. Plays a role in postnatal myogenesis, is involved in satellite cell activation (By similarity). Positively regulates IGF2 expression through PLAG1 and in a PLAG1-independent manner (PubMed:28796236).

Cellular Location


Nucleus.

HMGA2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HMGA2 Antibody (N-term) - Images



Western blot analysis of lysates from HepG2, NCCIT, mouse NIH/3T3 cell line (from left to right), using HMGA2 Antibody (N-term)(Cat. #AW5342). AW5342 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

HMGA2 Antibody (N-term) - Background

Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2. Plays an important role in chromosome condensation during the meiotic G2/M transition of spermatocytes.

HMGA2 Antibody (N-term) - References

- Patel U.A., et al. *Biochem. Biophys. Res. Commun.* 201:63-70(1994).
Schoenmakers E.F.P.M., et al. *Nat. Genet.* 10:436-444(1995).
Chau K.-Y., et al. *Nucleic Acids Res.* 23:4262-4266(1995).
Ashar H.R., et al. *Genomics* 31:207-214(1996).
Hauke S., et al. *Biochim. Biophys. Acta* 1729:24-31(2005).