

EPCAM Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5357

Specification

EPCAM Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P16422
Other Accession	Q75QW1 , Q3T0L5
Reactivity	Human
Predicted	Bovine, Pig
Host	Rabbit
Clonality	polyclonal
Calculated MW	H=35;M=35;Rat=35 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

EPCAM Antibody (C-term) - Additional Information

Gene ID 4072

Antigen Region
299-334

Other Names

Epithelial cell adhesion molecule, Ep-CAM, Adenocarcinoma-associated antigen, Cell surface glycoprotein Trop-1, Epithelial cell surface antigen, Epithelial glycoprotein, EGP, Epithelial glycoprotein 314, EGP314, hEGP314, KS 1/4 antigen, KSA, Major gastrointestinal tumor-associated protein GA733-2, T

Dilution

WB~~1:1000

Target/Specificity

This EPCAM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 299-334 amino acids from the C-terminal region of human EPCAM.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EPCAM Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EPCAM Antibody (C-term) - Protein Information

Name EPCAM

Synonyms GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO

Function

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

Cellular Location

Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction.
Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction

Tissue Location


Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

EPCAM Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EPCAM Antibody (C-term) - Images



All lanes : Anti-EPCAM Antibody (C-term)(AW5357) at 1/1000 dilution Lane 1: A431 whole cell lysates Lane 2: HT-29 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

EPCAM Antibody (C-term) - Background

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

EPCAM Antibody (C-term) - References

Strnad J., et al. Cancer Res. 49:314-317(1989).
Perez M.S., et al. J. Immunol. 142:3662-3667(1989).
Simon B., et al. Proc. Natl. Acad. Sci. U.S.A. 87:2755-2759(1990).
Szala S., et al. Proc. Natl. Acad. Sci. U.S.A. 87:3542-3546(1990).
Linnenbach A.J., et al. Mol. Cell. Biol. 13:1507-1515(1993).