

**EPOR Antibody (C-term)  
Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AW5363****Specification**

---

**EPOR Antibody (C-term) - Product Information**

Application	<b>WB, FC,E</b>
Primary Accession	<a href="#">P19235</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>H=55 KDa</b>
Isotype	<b>Rabbit IgG</b>
Antigen Source	<b>HUMAN</b>

**EPOR Antibody (C-term) - Additional Information****Gene ID** 2057**Antigen Region**  
470-504**Other Names**  
Erythropoietin receptor, EPO-R, EPOR**Dilution**  
WB~~1:1000  
FC~~1:25**Target/Specificity**  
This EPOR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 470-504 amino acids from the C-terminal region of human EPOR.**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.**Precautions**  
EPOR Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.**EPOR Antibody (C-term) - Protein Information****Name** EPOR

### Function

Receptor for erythropoietin. Mediates erythropoietin-induced erythroblast proliferation and differentiation. Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade. In some cell types, can also activate STAT1 and STAT3. May also activate the LYN tyrosine kinase.

### Cellular Location

Cell membrane; Single-pass type I membrane protein

### Tissue Location


Erythroid cells and erythroid progenitor cells. Isoform EPOR-F is the most abundant form in EPO-dependent erythroleukemia cells and in late-stage erythroid progenitors. Isoform EPOR-S and isoform EPOR-T are the predominant forms in bone marrow Isoform EPOR-T is the most abundant from in early-stage erythroid progenitor cells

## EPOR Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## EPOR Antibody (C-term) - Images




All lanes : Anti-EPOR Antibody (C-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 55 kDa Blocking/Dilution buffer: 5% NFD/MBST.



All lanes : Anti-EPOR Antibody (C-term)(AW5363) at 1/1000 dilution Lane 1: human kidney lysates  
 Lane 2: HL-60 whole cell lysates Lane 3: Jurkat whole cell lysates Lane 4: K562 whole cell lysates  
 Lane 5: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit  
 IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 62 kDa  
 Blocking/Dilution buffer: 5% NFD/MTBST.



Overlay histogram showing K562 cells stained with AW5363(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AW5363, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Flow cytometric analysis of K562 cells using EPOR Antibody (C-term)(green, Cat#AW5363) compared to an isotype control of rabbit IgG(blue). AW5363 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

#### **EPOR Antibody (C-term) - Background**

Receptor for erythropoietin. Mediates erythropoietin- induced erythroblast proliferation and differentiation. Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade. In some cell types, can also activate STAT1 and STAT3. May also activate the LYN tyrosine kinase.

#### **EPOR Antibody (C-term) - References**

- Winkelmann J.C.,et al.Blood 76:24-30(1990).
- Jones S.S.,et al.Blood 76:31-35(1990).
- Noguchi C.T.,et al.Blood 78:2548-2556(1991).
- Ehrenman K.,et al.Exp. Hematol. 19:973-977(1991).
- Nakamura Y.,et al.Science 257:1138-1141(1992).