

**CTSD(heavy chain) Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5367****Specification**

---

**CTSD(heavy chain) Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P07339</a>
Other Accession	<a href="#">P18242</a> , <a href="#">P80209</a>
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=45;M=45;Rat=45 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**CTSD(heavy chain) Antibody (Center) - Additional Information****Gene ID** 1509**Antigen Region**  
241-273**Other Names**

Cathepsin D, Cathepsin D light chain, Cathepsin D heavy chain, CTSD, CPSD

**Dilution**

WB~~1:1000

**Target/Specificity**

This CTSD(heavy chain) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 241-273 amino acids from the Central region of human CTSD(heavy chain).

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CTSD(heavy chain) Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**CTSD(heavy chain) Antibody (Center) - Protein Information**

**Name** CTSD

**Synonyms** CPSD

**Function**

Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed:<a href="http://www.uniprot.org/citations/27333034" target="\_blank">27333034</a>). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

**Cellular Location**

Lysosome. Melanosome. Secreted, extracellular space. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380)

**Tissue Location**


Expressed in the aorta extracellular space (at protein level) (PubMed:20551380). Expressed in liver (at protein level) (PubMed:1426530).

**CTSD(heavy chain) Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CTSD(heavy chain) Antibody (Center) - Images**



All lanes : Anti-CTSD(heavy chain) Antibody (Center) at 1/1000 dilution Lane 1: A431 whole cell lysates Lane 2: MCF-7 whole cell lysates Lane 3: SK-BR-3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution

Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### **CTSD(heavy chain) Antibody (Center) - Background**

Acid protease active in intracellular protein breakdown. Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

### **CTSD(heavy chain) Antibody (Center) - References**

Faust P.L.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:4910-4914(1985).

Westley B.R.,et al.Nucleic Acids Res. 15:3773-3786(1987).

Redecker B.,et al.DNA Cell Biol. 10:423-431(1991).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.