

UQCRQ Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5464

Specification

UQCRQ Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O14949
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=10;M=10;R=10 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

UQCRQ Antibody (N-term) - Additional Information

Gene ID 27089

Antigen Region
13-41

Other Names

Cytochrome b-c1 complex subunit 8, Complex III subunit 8, Complex III subunit VIII, Ubiquinol-cytochrome c reductase complex 95 kDa protein, Ubiquinol-cytochrome c reductase complex ubiquinone-binding protein QP-C, UQCRQ

Dilution

WB~~1:1000

Target/Specificity

This UQCRQ antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 13-41 amino acids from the N-terminal region of human UQCRQ.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UQCRQ Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UQCRQ Antibody (N-term) - Protein Information

Name UQCRQ**Function**

Component of the ubiquinol-cytochrome c oxidoreductase, a multisubunit transmembrane complex that is part of the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. The cytochrome b-c1 complex catalyzes electron transfer from ubiquinol to cytochrome c, linking this redox reaction to translocation of protons across the mitochondrial inner membrane, with protons being carried across the membrane as hydrogens on the quinol. In the process called Q cycle, 2 protons are consumed from the matrix, 4 protons are released into the intermembrane space and 2 electrons are passed to cytochrome c.


Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P08525}; Single-pass membrane protein {ECO:0000250|UniProtKB:P08525}

UQCRQ Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UQCRQ Antibody (N-term) - Images

All lanes : Anti-UQCRQ Antibody (N-term) at 1:1000 dilution Lane 1: human liver lysates Lane 2: HeLa whole cell lysates Lane 3: MCF-7 whole cell lysates Lane 4: NCI-H460 whole cell lysates

Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 10 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

UQCQRQ Antibody (N-term) - Background

This is a component of the ubiquinol-cytochrome c reductase complex (complex III or cytochrome b-c1 complex), which is part of the mitochondrial respiratory chain. This subunit, together with cytochrome b, binds to ubiquinone.

UQCQRQ Antibody (N-term) - References

Fujiwara T., et al. Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
Schaeffer H., et al. Methods Enzymol. 260:82-96(1995).
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).
Barel O., et al. Am. J. Hum. Genet. 82:1211-1216(2008).