

CDA Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5473

Specification

CDA Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P32320
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=16;M=16;R=16 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

CDA Antibody (Center) - Additional Information

Gene ID 978

Antigen Region
51-83

Other Names
Cytidine deaminase, Cytidine aminohydrolase, CDA, CDD

Dilution
WB~~1:1000

Target/Specificity
This CDA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 51-83 amino acids from the Central region of human CDA.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CDA Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CDA Antibody (Center) - Protein Information

Name CDA ([HGNC:1712](#))

Synonyms CDD

Function

This enzyme scavenges exogenous and endogenous cytidine and 2'-deoxycytidine for UMP synthesis.

Tissue Location


Highly expressed in granulocytes while expression is very low in fibroblasts, chondrocytes, monocytes, and T- as well as B-cell lines

CDA Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDA Antibody (Center) - Images



All lanes : Anti-CDA Antibody (Center) at 1:1000 dilution Lane 1: human liver lysates Lane 2: HeLa whole cell lysates Lane 3: mouse kidney lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 16 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

CDA Antibody (Center) - Background

CDA forms a homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. This protein is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in the CDA gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine arabinoside used in the treatment of certain childhood leukemias.

CDA Antibody (Center) - References

Laliberte J., Momparler R.L. *Cancer Res.* 54:5401-5407(1994)

Demontis S., Terao M. *Biochim. Biophys. Acta* 1443:323-333(1998)

Kuhn K., Bertling W.M., Emmrich F. *Biochem. Biophys. Res. Commun.* 190:1-7(1993)