

**LEFTY2 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5566**

**Specification**

---

**LEFTY2 Antibody - Product Information**

Application	WB,E
Primary Accession	<a href="#">O00292</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=41,37;M=41 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**LEFTY2 Antibody - Additional Information**

**Gene ID** 7044

**Antigen Region**  
1-366

**Other Names**

Left-right determination factor 2, Endometrial bleeding-associated factor, Left-right determination factor A, Protein lefty-2, Protein lefty-A, Transforming growth factor beta-4, TGF-beta-4, LEFTY2, EBAF, LEFTA, LEFTYA, TGFB4

**Dilution**

WB~~1:8000

**Target/Specificity**

This LEFTY2 antibody is generated from a rabbit immunized with a recombinant.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

LEFTY2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**LEFTY2 Antibody - Protein Information**

**Name** LEFTY2

**Synonyms** EBAF, LEFTA, LEFTYA, TGFB4

**Function**

Required for left-right (L-R) asymmetry determination of organ systems in mammals. May play a

role in endometrial bleeding.

#### Cellular Location

Secreted.

#### Tissue Location


Mesenchymal cells of the endometrial stroma.

### LEFTY2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### LEFTY2 Antibody - Images



All lanes : Anti-LEFTY2 Antibody at 1:8000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

### LEFTY2 Antibody - Background

Required for left-right (L-R) asymmetry determination of organ systems in mammals. May play a role in endometrial bleeding.

### LEFTY2 Antibody - References

Kothapalli R., et al. J. Clin. Invest. 99:2342-2350(1997).  
Kosaki K., et al. Am. J. Hum. Genet. 64:712-721(1999).

Ota T., et al. Nat. Genet. 36:40-45(2004).  
Otsuki T., et al. DNA Res. 12:117-126(2005).  
Gregory S.G., et al. Nature 441:315-321(2006).