

**DBNL Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5587**

**Specification**

---

**DBNL Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9UJU6</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=48,49,37,43,38 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**DBNL Antibody (C-term) - Additional Information**

**Gene ID** 28988

**Antigen Region**  
292-319

**Other Names**

Drebrin-like protein, Cervical SH3P7, Cervical mucin-associated protein, Drebrin-F, HPK1-interacting protein of 55 kDa, HIP-55, SH3 domain-containing protein 7, DBNL, CMAP, SH3P7

**Dilution**

WB~~1:2000

**Target/Specificity**

This DBNL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 292-319 amino acids from the C-terminal region of human DBNL.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DBNL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**DBNL Antibody (C-term) - Protein Information**

**Name** DBNL

**Synonyms** CMAP, SH3P7

**Function**

Adapter protein that binds F-actin and DNM1, and thereby plays a role in receptor-mediated endocytosis. Plays a role in the reorganization of the actin cytoskeleton, formation of cell projections, such as neurites, in neuron morphogenesis and synapse formation via its interaction with WASL and COBL. Does not bind G-actin and promote actin polymerization by itself. Required for the formation of organized podosome rosettes (By similarity). May act as a common effector of antigen receptor-signaling pathways in leukocytes. Acts as a key component of the immunological synapse that regulates T-cell activation by bridging TCRs and the actin cytoskeleton to gene activation and endocytic processes.

#### Cellular Location

Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q62418}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q62418}. Cell projection, ruffle {ECO:0000250|UniProtKB:Q62418}. Cytoplasm, cell cortex {ECO:0000250|UniProtKB:Q62418}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9JHL4}. Synapse {ECO:0000250|UniProtKB:Q62418} Perikaryon {ECO:0000250|UniProtKB:Q62418}. Cell projection, neuron projection {ECO:0000250|UniProtKB:Q62418}. Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Cytoplasmic vesicle, clathrin-coated vesicle membrane {ECO:0000250|UniProtKB:Q62418}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q62418}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Cell projection, podosome {ECO:0000250|UniProtKB:Q62418}. Early endosome. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9JHL4}. Postsynaptic density {ECO:0000250|UniProtKB:Q9JHL4}. Note=Associates with lamellipodial actin and membrane ruffles. Colocalizes with actin and cortactin at podosome dots and podosome rosettes. {ECO:0000250|UniProtKB:Q62418, ECO:0000250|UniProtKB:Q9JHL4}


#### DBNL Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### DBNL Antibody (C-term) - Images





All lanes : Anti-DBNL Antibody (C-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: SH-SY5Y whole cell lysate Lane 4: MOLT-4 whole cell lysate Lane 5: A431 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

#### **DBNL Antibody (C-term) - Background**

Actin-binding adapter protein. May act as a common effector of antigen receptor-signaling pathways in leukocytes. Its association with dynamin suggests that it may also connect the actin cytoskeleton to endocytic function. Acts as a key component of the immunological synapse that regulates T-cell activation by bridging TCRs and the actin cytoskeleton to gene activation and endocytic processes. Binds to F-actin but is not involved in actin polymerization, capping or bundling. Does not bind G-actin.

#### **DBNL Antibody (C-term) - References**

Venkatesan, K., et al. Nat. Methods 6(1):83-90(2009)  
Haeckel, A., et al. J. Neurosci. 28(40):10031-10044(2008)  
Le Bras, S., et al. FEBS Lett. 581(5):967-974(2007)  
Lamesch, P., et al. Genomics 89(3):307-315(2007)  
Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :