

**MFRN2 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5588**

**Specification**

---

**MFRN2 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O96A46</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=39 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**MFRN2 Antibody (N-term) - Additional Information**

**Gene ID** 81894

**Antigen Region**  
36-64

**Other Names**

Mitoferrin-2, Mitochondrial RNA-splicing protein 3/4 homolog, MRS3/4, hMRS3/4, Mitochondrial iron transporter 2, Solute carrier family 25 member 28, SLC25A28, MFRN2

**Dilution**

WB~~1:2000  
IHC-P~~1:50~100  
FC~~1:10~50

**Target/Specificity**

This MFRN2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 36-64 amino acids from the N-terminal region of human MFRN2.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MFRN2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**MFRN2 Antibody (N-term) - Protein Information**

**Name** SLC25A28

**Synonyms** MFRN2

### Function

Mitochondrial iron transporter that mediates iron uptake. Probably required for heme synthesis of hemoproteins and Fe-S cluster assembly in non-erythroid cells.

### Cellular Location

[Isoform 1]: Mitochondrion inner membrane; Multi-pass membrane protein

### Tissue Location


Ubiquitous. Expressed in placenta, lung, kidney, pancreas, liver, brain, skeletal muscle and heart

## MFRN2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## MFRN2 Antibody (N-term) - Images



All lanes : Anti-MFRN2 Antibody (N-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



MFRN2 Antibody (N-term) (Cat. #AW5588) IHC analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MFRN2 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



MFRN2 Antibody (N-term) (Cat. #AW5588) flow cytometric analysis of Jurkat cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **MFRN2 Antibody (N-term) - Background**

MFRN2 is mitochondrial iron transporter that mediates iron uptake. It is probably required for heme synthesis of hemoproteins and Fe-S cluster assembly in non-erythroid cells. The iron delivered into the mitochondria, presumably as Fe(2+), is then probably delivered to ferrochelatase to catalyze Fe(2+) incorporation into protoporphyrin IX to make heme.

### **MFRN2 Antibody (N-term) - References**

Grupe, A., et al. *Am. J. Hum. Genet.* 78(1):78-88(2006)  
Deloukas, P., et al. *Nature* 429(6990):375-381(2004)  
Wistow, G., et al. *Mol. Vis.* 8, 185-195 (2002)