

NDUFV1 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5622

Specification

NDUFV1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P49821
Other Accession	P25708 , Q54190 , Q0MQI5 , Q8HXQ9 , Q91YT0 , Q0MQI6 , Q0MQI4
Reactivity	Human, Mouse
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=51,50;M=51 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

NDUFV1 Antibody (Center) - Additional Information

Gene ID 4723

Antigen Region
194-226

Other Names

NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial, Complex I-51kD, CI-51kD, NADH dehydrogenase flavoprotein 1, NADH-ubiquinone oxidoreductase 51 kDa subunit, NDUFV1, UQOR1

Dilution
WB~~0.25

Target/Specificity

This NDUFV1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 194-226 amino acids from the Central region of human NDUFV1.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NDUFV1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NDUFV1 Antibody (Center) - Protein Information

Name NDUFV1 ([HGNC:7716](#))

Synonyms UQOR1

Function

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:28844695). Part of the peripheral arm of the enzyme, where the electrons from NADH are accepted by flavin mononucleotide (FMN) and then passed along a chain of iron-sulfur clusters by electron tunnelling to the final acceptor ubiquinone (PubMed:28844695). Contains FMN, which is the initial electron acceptor as well as one iron-sulfur cluster (PubMed:28844695).

Cellular Location


Mitochondrion inner membrane {ECO:0000250|UniProtKB:P25708}; Peripheral membrane protein {ECO:0000250|UniProtKB:P25708}; Matrix side {ECO:0000250|UniProtKB:P25708}

NDUFV1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDUFV1 Antibody (Center) - Images



All lanes : Anti-NDUFV1 Antibody (Center) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: mouse kidney lysate Lane 5: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

NDUFV1 Antibody (Center) - Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

NDUFV1 Antibody (Center) - References

de Coo R.F.M.,et al.Mamm. Genome 10:49-53(1999).
Schuelke M.,et al.Biochem. Biophys. Res. Commun. 245:599-606(1998).
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.