

DANRE ywhag1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb10022c

Specification

DANRE ywhag1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q6PC29
Reactivity	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28235
Antigen Region	137-162

DANRE ywhag1 Antibody (Center) - Additional Information

Gene ID 117604

Other Names

14-3-3 protein gamma-1, ywhag1, ywhag

Target/Specificity

This DANRE ywhag1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 137-162 amino acids from the Central region of DANRE ywhag1.

Dilution

WB~~1:8000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DANRE ywhag1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DANRE ywhag1 Antibody (Center) - Protein Information

Name ywhag1

Synonyms ywhag

Function Adapter protein implicated in the regulation of a large spectrum of both general and

specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner.

Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:P68252}.

DANRE ywhag1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DANRE ywhag1 Antibody (Center) - Images



DANRE ywhag1 Antibody (Center) (Cat. #Azb10022c) western blot analysis in zebra fish brain tissue lysates (35ug/lane). This demonstrates the DANRE ywhag1 antibody detected the DANRE ywhag1 protein (arrow).



Anti-DANRE ywhag1 Antibody (Center) at 1:8000 dilution + ZF4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

DANRE ywhag1 Antibody (Center) - Background

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner (By similarity).