

ACTR3 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP2891b**Specification**

ACTR3 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P61158](#)**ACTR3 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 10096

Other Names

Actin-related protein 3, Actin-like protein 3, ACTR3, ARP3

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP2891b](/products/AP2891b) was selected from the C-term region of human ACTR3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ACTR3 Antibody (C-term) Blocking Peptide - Protein Information

Name ACTR3

Synonyms ARP3

Function

ATP-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF) (PubMed:[9000076](http://www.uniprot.org/citations/9000076)). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility (PubMed:[9000076](http://www.uniprot.org/citations/9000076)). Seems to contact the pointed end of the daughter actin filament (PubMed:[9000076](http://www.uniprot.org/citations/9000076)). In podocytes, required for the formation of lamellipodia downstream of AVIL and PLCE1 regulation (PubMed:[29058690](http://www.uniprot.org/citations/29058690)). In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin

polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:17220302, PubMed:29925947). The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed:29925947). Plays a role in ciliogenesis (PubMed:20393563).

Cellular Location

Cytoplasm, cytoskeleton. Cell projection. Nucleus. Note=In pre- apoptotic cells, colocalizes with MEFV in large specks (pyroptosomes) (PubMed:19109554)

ACTR3 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ACTR3 Antibody (C-term) Blocking Peptide - Images

ACTR3 Antibody (C-term) Blocking Peptide - Background

ACTR3 is known to be a major constituent of the ARP2/3 complex. This complex is located at the cell surface and is essential to cell shape and motility through lamellipodial actin assembly and protrusion.

ACTR3 Antibody (C-term) Blocking Peptide - References

Weisswange,I., et. al., Nature 458 (7234), 87-91 (2009)