

**ALKBH1 Antibody (C-term) Blocking peptide**  
Synthetic peptide  
Catalog # BP5867b**Specification**

---

**ALKBH1 Antibody (C-term) Blocking peptide - Product Information**

Primary Accession [O13686](#)  
Other Accession [NP\\_006011.2](#)

**ALKBH1 Antibody (C-term) Blocking peptide - Additional Information**

Gene ID 8846

**Other Names**

Alkylated DNA repair protein alkB homolog 1, Alpha-ketoglutarate-dependent dioxygenase ABH1, DNA lyase ABH1, DNA oxidative demethylase ALKBH1, ALKBH1, ABH, ABH1, ALKBH

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**ALKBH1 Antibody (C-term) Blocking peptide - Protein Information**

Name ALKBH1 ([HGNC:17911](#))

**Function**

Dioxygenase that acts as on nucleic acids, such as DNA and tRNA (PubMed:[18603530](http://www.uniprot.org/citations/18603530), PubMed:[27497299](http://www.uniprot.org/citations/27497299), PubMed:[27745969](http://www.uniprot.org/citations/27745969)). Requires molecular oxygen, alpha-ketoglutarate and iron (PubMed:[18603530](http://www.uniprot.org/citations/18603530), PubMed:[27497299](http://www.uniprot.org/citations/27497299)). A number of activities have been described for this dioxygenase, but recent results suggest that it mainly acts as on tRNAs and mediates their demethylation or oxidation depending on the context and subcellular compartment (PubMed:[27497299](http://www.uniprot.org/citations/27497299), PubMed:[27745969](http://www.uniprot.org/citations/27745969)). Mainly acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs, with a preference for N(1)-methyladenine at position 58 (m1A58) present on a stem loop structure of tRNAs (PubMed:[27745969](http://www.uniprot.org/citations/27745969)). Acts as a regulator of translation initiation and elongation in response to glucose deprivation: regulates both

translation initiation, by mediating demethylation of tRNA(Met), and translation elongation, N(1)-methyladenine-containing tRNAs being preferentially recruited to polysomes to promote translation elongation (PubMed:[27745969](http://www.uniprot.org/citations/27745969)). In mitochondrion, specifically interacts with mt-tRNA(Met) and mediates oxidation of mt-tRNA(Met) methylated at cytosine(34) to form 5- formylcytosine (f(5)c) at this position (PubMed:[27497299](http://www.uniprot.org/citations/27497299)). mt-tRNA(Met) containing the f(5)c modification at the wobble position enables recognition of the AUA codon in addition to the AUG codon, expanding codon recognition in mitochondrial translation (PubMed:[27497299](http://www.uniprot.org/citations/27497299)). Specifically demethylates DNA methylated on the 6th position of adenine (N(6)-methyladenosine) DNA (PubMed:[30017583](http://www.uniprot.org/citations/30017583)), PubMed:[30392959](http://www.uniprot.org/citations/30392959)). N(6)- methyladenosine (m6A) DNA is present at some L1 elements in embryonic stem cells and probably promotes their silencing (By similarity). Demethylates mRNAs containing N(3)-methylcytidine modification (PubMed:[31188562](http://www.uniprot.org/citations/31188562)). Also able to repair alkylated single-stranded DNA by oxidative demethylation, but with low activity (PubMed:[18603530](http://www.uniprot.org/citations/18603530)). Also has DNA lyase activity and introduces double-stranded breaks at abasic sites: cleaves both single-stranded DNA and double-stranded DNA at abasic sites, with the greatest activity towards double-stranded DNA with two abasic sites (PubMed:[19959401](http://www.uniprot.org/citations/19959401)). DNA lyase activity does not require alpha-ketoglutarate and iron and leads to the formation of an irreversible covalent protein-DNA adduct with the 5' DNA product (PubMed:[19959401](http://www.uniprot.org/citations/19959401)), PubMed:[23577621](http://www.uniprot.org/citations/23577621)). DNA lyase activity is not required during base excision repair and class switch recombination of the immunoglobulin heavy chain during B lymphocyte activation. May play a role in placental trophoblast lineage differentiation (By similarity).

#### **Cellular Location**

Nucleus. Mitochondrion. Note=Mainly localizes in euchromatin, largely excluded from heterochromatin and nucleoli (By similarity). {ECO:0000250|UniProtKB:POCB42}

#### **Tissue Location**

Ubiquitous.

### **ALKBH1 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **ALKBH1 Antibody (C-term) Blocking peptide - Images**