

## **AGXT Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP6500c

## **Specification**

## **AGXT Antibody (Center) Blocking Peptide - Product Information**

Primary Accession

P21549

# **AGXT Antibody (Center) Blocking Peptide - Additional Information**

Gene ID 189

#### **Other Names**

Serine--pyruvate aminotransferase, SPT, Alanine--glyoxylate aminotransferase, AGT, AGXT, AGT1, SPAT

## Target/Specificity

The synthetic peptide sequence used to generate the antibody <a

href=/products/AP6500c>AP6500c</a> was selected from the Center region of human AGXT. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### AGXT Antibody (Center) Blocking Peptide - Protein Information

Name AGXT (HGNC:341)

Synonyms AGT1, SPAT

## **Function**

Peroxisomal aminotransferase that catalyzes the transamination of glyoxylate to glycine and contributes to the glyoxylate detoxification (PubMed:<a

href="http://www.uniprot.org/citations/10960483" target="\_blank">10960483</a>, PubMed:<a href="http://www.uniprot.org/citations/12777626" target="\_blank">12777626</a>, PubMed:<a href="http://www.uniprot.org/citations/23229545" target="\_blank">23229545</a>, PubMed:<a href="http://www.uniprot.org/citations/24055001" target="\_blank">24055001</a>, PubMed:<a href="http://www.uniprot.org/citations/26149463" target="\_blank">26149463</a>). Also catalyzes the transamination between L-serine and pyruvate and contributes to gluconeogenesis from the L-serine metabolism (PubMed:<a href="http://www.uniprot.org/citations/10347152"



target="\_blank">10347152</a>).

**Cellular Location**Peroxisome

Tissue Location Liver.

# **AGXT Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

**AGXT Antibody (Center) Blocking Peptide - Images** 

AGXT Antibody (Center) Blocking Peptide - Background

AGXT is expressed only in the liver and protein is localized mostly in the peroxisomes, where it is involved in glyoxylate detoxification.

# **AGXT Antibody (Center) Blocking Peptide - References**

Cellini,B., J. Biol. Chem. 284 (13), 8349-8358 (2009)Bertoldi,M., Biochim. Biophys. Acta 1784 (9), 1356-1362 (2008)