

**ALB Antibody (C-term) Blocking Peptide**  
Synthetic peptide  
Catalog # BP6540b**Specification**

---

**ALB Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P02768](#)**ALB Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 213

**Other Names**

Serum albumin, ALB

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6540b](/products/AP6540b) was selected from the C-term region of human ALB. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**ALB Antibody (C-term) Blocking Peptide - Protein Information**

Name ALB

**Function**

Binds water, Ca(2+), Na(+), K(+), fatty acids, hormones, bilirubin and drugs (Probable). Its main function is the regulation of the colloidal osmotic pressure of blood (Probable). Major zinc transporter in plasma, typically binds about 80% of all plasma zinc (PubMed:[19021548](http://www.uniprot.org/citations/19021548)). Major calcium and magnesium transporter in plasma, binds approximately 45% of circulating calcium and magnesium in plasma (By similarity). Potentially has more than two calcium-binding sites and might additionally bind calcium in a non-specific manner (By similarity). The shared binding site between zinc and calcium at residue Asp-273 suggests a crosstalk between zinc and calcium transport in the blood (By similarity). The rank order of affinity is zinc > calcium > magnesium (By similarity). Binds to the bacterial siderophore enterobactin and inhibits enterobactin-mediated iron uptake of E.coli from ferric transferrin, and may thereby limit the utilization of iron and growth of enteric bacteria such as E.coli (PubMed:[6234017](http://www.uniprot.org/citations/6234017))

target="\_blank">6234017</a>). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>).

**Cellular Location**

Secreted.

**Tissue Location**

Plasma.

**ALB Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**ALB Antibody (C-term) Blocking Peptide - Images****ALB Antibody (C-term) Blocking Peptide - Background**

Albumin is a soluble, monomeric protein which comprises about one-half of the blood serum protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. Albumin is synthesized in the liver as prealbumin which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted albumin.

**ALB Antibody (C-term) Blocking Peptide - References**

Liu,Z., Biophys. J. 96 (10), 3917-3925 (2009)Minchiotti,L., Hum. Mutat. 29 (8), 1007-1016 (2008)